



RESEARCH

Open Access

The impact of HIV/AIDS on compliance with antidepressant treatment in major depressive disorder: A prospective study in a South African private healthcare cohort

Francois N Slabbert¹, Brian H Harvey^{2,3}, Christiaan B Brink² and Martie S Lubbe^{1*}

Abstract

Background: MDD and HIV/AIDS have a high prevalence worldwide with severe consequences for patients. In both conditions, compliance with treatment is key to successfully treat these disorders. In the current study, we examine the effect of MDD on the compliance with ADs in patients diagnosed with co-morbid HIV/AIDS and how different classes of ADs influence compliance in this group of patients.

Methods: A prospective, cohort study design was used to analyse nationally representative medicine claims data submitted to a privately-owned South African Pharmaceutical Benefit Management (PBM) company. Two groups were distinguished in the database, namely patients with only MDD and patients with both MDD and HIV/AIDS, over a six-year study period. The study population was determined by the following inclusion criteria: patients older than 18 years, MDD should be diagnosed by a psychiatrist supported by an appropriate ICD-10 code, and all patients have to be on combination antiretroviral treatment (cARV) treatment. The medicine possession ratio (MPR) was used as proxy to determine patient compliance with AD medication.

Results: 127 patients (i.e. 0.24%) met the criteria of co-morbid MDD and HIV/AIDS. Females have a significantly higher prevalence of MDD and HIV/AIDS when compared to males. Patients diagnosed with both HIV/AIDS and MDD (74.43 ± 32.03 , 95% CI: 71.51-77.34) have a statistical significantly ($p < 0.0001$) lower compliance with AD treatment vs. MDD patients ($80.94\% \pm 29.44$, 95% CI: 80.56-81.33), but the practical significance thereof, is low (Cohen's $d = 0.2255$). In this group only 26.83% of TCA had acceptable compliance compared to the 58.57% of SNRIs. Noteworthy observations were that 75% ($p < 0.0217$; Cramer's $V = 0.0388$) of venlafaxine and 28.6% ($p < 0.0197$; Cramer's $V = -0.0705$) of the paroxetine items were compliant in patients diagnosed with both HIV/AIDS and MDD.

Conclusions: AD compliance is statistical significantly lower in depressed HIV/AIDS vs. depressed non-HIV/AIDS patients. However, these differences is of low practical or clinical significance, meaning that depressed HIV/AIDS patients would have missed approximately two AD doses (6.5% of a 30-day treatment period) more than the non-HIV/AIDS depressed patient over the same treatment period.

Keywords: Major depressive disorder, HIV/AIDS, Compliance, Antidepressants, Venlafaxine

* Correspondence: Martie.Lubbe@nwu.ac.za

¹Medicines Usage Group (MUSA), North-West University, Potchefstroom 2520, South Africa

Full list of author information is available at the end of the article

Background

In 2012, the UNAIDS World AIDS Day Report estimated that approximately 34 million people worldwide are infected with human immunodeficiency virus (HIV), whereas 69% of these infected patients live in sub-Saharan Africa [1]. In the United States, more than one million individuals are infected with HIV and acquired immunodeficiency syndrome (AIDS, collectively HIV/AIDS), while nearly 50 000 more contract HIV each year [2]. In South Africa, the prevalence of HIV/AIDS is estimated to be 17.3% in the age group 15 to 49 years and approximately 5.6 million people in the country are living with HIV/AIDS [3]. South Africa is one of the countries with the highest HIV/AIDS-related mortality rates in the world, with approximately 270 000 deaths reported in 2011 [3].

Apart from the huge impact on the health system, HIV/AIDS also affects individual patients, causing a great deal of suffering in both their social lives as well as mental and physical health. It is therefore not surprising that HIV/AIDS is co-morbid with various psychosocial disorders, such as major depressive disorder (MDD) [4]. MDD is one of the main causes of psychiatric morbidity in HIV/AIDS patients [4]. In a six-month follow-up study in HIV/AIDS-infected patients in South Africa, Olley and co-workers described a 34.9% prevalence of MDD [5]. Elsewhere in the literature, however, there is some controversy over the prevalence of MDD in HIV/AIDS patients, with the prevalence rate varying between 3 and 54% [4,6,7]. Further studies in this regard are therefore urgently needed.

The co-morbidity between chronic illnesses and MDD is a well-recognised phenomenon. Chronically ill patients display two to three times higher rates of MDD when compared to healthy individuals of the same age and gender [8-10]. Diabetes, asthma, epilepsy, cancer, coronary heart disease, hypertension and HIV/AIDS are among the most common chronic illnesses that are associated with an increased prevalence of MDD [11,12]. Considering the symptomatology of depression, depressed patients tend to neglect themselves due to fatigue, low energy levels, impaired memory and the sense of helplessness [13]. Patients with a chronic illness and co-morbid MDD have a significantly decreased compliance with self-care treatment regimens [14], which negatively affects long-term treatment outcomes and is ultimately associated with impaired daily functioning and a decreased quality of life [15,16].

HIV-positive patients have twice the incidence of MDD compared to HIV-negative individuals [17-20]. In particular, late stage HIV (CD4 count of <200 cells/ μ L) is associated with a significantly higher incidence of MDD, with a number of studies showing as much as a two and a half-fold increase in the incidence of MDD

[21-24]. A recent study by Lopes and co-workers found that HIV-positive men were more likely to have a major depressive disorder/dysthymia (OR = 3.77; 95% CI, 1.16-12.27) when compared to HIV-negative males [25]. The reason for this is likely to be multifactorial, although chronic-psychosocial stress (e.g. discrimination, isolation, violence, stigmatisation, hopelessness and drug abuse) [26] associated with HIV/AIDS can induce inflammation in the central nervous system (CNS), which is a major etiological factor [27]. Exposure to chronic psychosocial stress induces the continuous release of cortisol, and activates the hypothalamic-pituitary-adrenal (HPA) axis as well as the sympathetic autonomic nervous system, which in turn activates immune cells in both CNS and in the periphery [28,29]. Depressed patients express elevated levels and an imbalance of cytokines, especially interleukin-1 (IL-1) and tumour necrosis factor alpha (TNF α), which leads to cognitive and mood impairment [30-32]. Further evidence supporting inflammation as an etiological factor in depression, is that patients receiving interferon Alfa (IFN- α) therapy for various types of cancer develop severe depression, INF being an inflammatory mediator [33-35]. Cerebral HIV infection will induce chronic inflammation in the brain and this causes the release of IFN and other cytokines, with depression as a result [27]. Therefore, HIV/AIDS patients with chronic increased cytokine levels as a result of viral infection, can be expected to be particularly vulnerable to develop depression [36].

The introduction of combination antiretroviral therapy (cART) has been successful in improving morbidity and decreasing the mortality rate of HIV/AIDS, which provides a glimmer of hope [37-39]. Importantly, however, cART can suppress HIV-1 RNA levels and increase CD4 T cell lymphocytes only if the cART is associated with excellent compliance and persistence [40-43]. Patients who are HIV positive must maintain a compliance rate of at least 95% in order to prevent virologic failure, increase CD4 T cell lymphocyte count, and decrease viral load and opportunistic infections [44-47]. Another risk associated with non-compliance with ART is the increased risk of viral resistance that ultimately leads to treatment failure [48-51]. On the other hand, compliance with AD treatment is just as important. Patients who fail to comply with their AD treatment have a two-fold increased risk for relapse [52], are more likely to experience antidepressant withdrawal syndrome [53], and have a significantly increased mortality and morbidity rate associated with MDD [54]. Consequently, poor compliance with either ART or AD treatment will significantly compromise a successful treatment outcome.

The current literature regarding the prevalence of HIV/AIDS and MDD is quite limited. Therefore, this study will firstly strive to determine the prevalence of

HIV/AIDS-positive patients within the MDD-diagnosed population in the private health sector of South Africa. Secondly, the study will investigate how AD compliance is affected by MDD in HIV/AIDS-positive patients when compared to depressed non-HIV/AIDS patients and, lastly, whether AD compliance has any association with gender and antidepressant class in this population.

Results

The research cohort consisted of two groups, namely MDD patients with HIV/AIDS (n = 127) and patients with only MDD (n = 12 270) (Table 1). These patients received over the study period respectively 466 and 22 831 AD items (Table 2).

The mean MPR of antidepressants (ADs) was statistically significantly lower ($p < 0.0001$; $d = 0.2255$) in MDD patients with HIV/AIDS ($74.43\% \pm 32.03$, 95% CI: 71.51-77.34) than in patients with only MDD ($80.94\% \pm 29.44$, 95% CI: 80.56-81.33). These differences were of low practical significance. In MDD patients with HIV/AIDS, antidepressant compliance (MPR) was statistical significantly lower with medium practical significance in females ($69.35\% \pm 29.69$, 95% CI: 66.03-72.67) than in males ($84.51\% \pm 34.18$, 95% CI: 79.10-89.91) from the same cohort ($p < 0.0001$; $d = 0.4755$). Age was not a significant predictor of AD compliance in MDD patients with HIV/AIDS ($p < 0.9519$) and those with only MDD ($p = 0.1111$), as shown in Table 2.

In Table 3, a significant association was found between AD class and AD compliance in MDD patients with HIV/AIDS, with compliance figures as follows; tricyclic antidepressants (TCA) 26.83%; selective serotonin re-

uptake inhibitors (SSRIs) 44.93%; diverse antidepressants 52.70%; and serotonin and noradrenalin re-uptake inhibitors (SNRIs) 58.57% ($p < 0.0001$; Cramer's $V = 0.3450$).

The following ADs, as depicted in Table 4, were associated with unacceptably low levels of compliance in MDD patients with HIV/AIDS vs. patients with only MDD, as observed from the compliance ratio: amitriptyline 23.53% vs. 39.34% ($p < 0.0618$; Cramer's $V = -0.0494$); trazodone 27.27% vs. 50.72% ($p < -0.0239$; Cramer's $V = -0.0630$); paroxetine 28.57% vs. 54.19% ($p < 0.0197$; Cramer's $V = -0.0705$); duloxetine 36.67% vs. 55.51% ($p < 0.0391$; Cramer's $V = -0.0426$); escitalopram 44.16% vs. 52.74% ($p < 0.0359$; Cramer's $V = -0.0254$); bupropion 78.26% vs. 50.46% ($p < 0.0082$; Cramer's $V = 0.0698$) and venlafaxine 75.00% vs. 56.93% ($p < 0.0217$; Cramer's $V = 0.0388$).

Discussion

This study focused on the prevalence of HIV/AIDS-positive patients within the MDD-diagnosed population in this section of the private health sector of South Africa. Secondly, the study investigated how AD compliance is affected by MDD in HIV/AIDS-positive patients when compared to depressed non-HIV/AIDS patients, and whether AD compliance has any correlation with gender and antidepressant class in this population. These goals were attained in that we were able to confirm the co-morbidity between MDD and HIV/AIDS-positive patients; however, due to the strict inclusion criteria, the numbers of patients were small, which provided a limited picture of the overall prevalence between MDD and HIV/AIDS-positive patients.

This study found that patients diagnosed with both HIV/AIDS and MDD (74.43 ± 32.03 , 95% CI: 71.51-77.34) have a statistical significantly lower compliance with AD treatment when compared to patients diagnosed only with MDD ($80.94\% \pm 29.44$, 95% CI: 80.56-81.33). However, these differences is of low practical or clinical significance, meaning that depressed HIV/AIDS patients would have missed approximately two AD doses (~6.5% of a 30-day treatment period) more than the non-HIV/AIDS depressed patient over the same treatment period. The current study confirms that patients suffering from both MDD and HIV/AIDS have a decreased compliance with MDD treatment. This trend is confirmed in the literature as several authors have found similar results, namely that patients suffering from both conditions are less compliant to AD treatment regimens [14,20,55,56]. Our data also seem to suggest, in Table 4, that patients with HIV/AIDS are less compliant with ADs that present with multi-receptor pharmacology, i.e. TCAs (26.83% compliance; Table 3) and paroxetine (compliance 28.57%, Table 4), while they are more compliant with pharmacologically 'clean' ADs, such as the venlafaxine (75% compliance, Table 4). The TCAs display a

Table 1 Patient demographics

| Total number of patients in the study population | | | | | | |
|--|----------------|-------|----------------|-------|-----------|-------|
| Illness | n | % | | | | |
| MDD + HIV | 127 | 0.24 | | | | |
| MDD | 12270 | 22.94 | | | | |
| HIV | 41086 | 76.82 | | | | |
| Gender classification of the study population | | | | | | |
| Illness | Male | | Female | | | |
| | n | % | n | % | | |
| MDD + HIV | 37 | 29.13 | 90 | 70.87 | | |
| MDD | 3496 | 28.49 | 8774 | 71.51 | | |
| HIV | 18268 | 44.46 | 22818 | 55.54 | | |
| Age classification of the study population | | | | | | |
| Illness | >18 ≤ 40 years | | >40 ≤ 60 years | | >60 years | |
| | n | % | n | % | n | % |
| MDD + HIV | 31 | 24.41 | 75 | 59.06 | 21 | 16.54 |
| MDD | 2662 | 21.7 | 21176 | 47.03 | 3837 | 31.27 |
| HIV | 16586 | 40.37 | 21176 | 51.54 | 1652 | 4.02 |

Table 2 The effect of illness, gender and age on medicine possession ratio (MPR)

| Mean MPR of ADs in MDD patients with/without HIV/AIDS (dispensed items) | | | | | |
|--|-------|---------------|-------------|--------|-----------------|
| Illness | n | Mean ± SD (%) | 95% CI | p* | Cohen's d-value |
| HIV/AIDS + MDD | 466 | 74.43 ± 32.03 | 71.51-77.34 | 0.0001 | 0.2255 |
| MDD | 22831 | 80.94 ± 29.44 | 80.56-81.33 | | |
| Mean MPR of ADs in MDD patients with HIV/AIDS vs. gender (dispensed items) | | | | | |
| Gender | n | Mean ± SD (%) | 95% CI | p* | Cohen's d-value |
| Female | 310 | 69.35 ± 29.69 | 66.03-72.67 | 0.0001 | 0.4755 |
| Male | 156 | 84.51 ± 34.18 | 79.10-89.91 | | |
| Mean MPR of ADs in MDD patients without HIV/AIDS vs. gender (dispensed items) | | | | | |
| Gender | n | Mean ± SD (%) | 95% CI | p* | |
| Female | 16600 | 80.97 ± 29.72 | 80.52-81.43 | 0.8067 | |
| Male | 6231 | 80.87 ± 28.65 | 80.16-81.58 | | |
| Mean MPR of ADs in MDD patients with HIV/AIDS vs. age group (dispensed items) | | | | | |
| Age (years) | n | Mean ± SD (%) | 95% CI | p** | |
| > 18 ≤ 40 | 126 | 74.57 ± 32.39 | 68.86-80.29 | 0.9519 | |
| >40 ≤ 60 | 284 | 74.61 ± 33.07 | 70.74-78.47 | | |
| >60 | 56 | 73.16 ± 25.74 | 66.27-80.06 | | |
| Mean MPR of ADs in MDD patients without HIV/AIDS vs. age group (dispensed items) | | | | | |
| Age (years) | n | Mean ± SD (%) | 95% CI | p** | |
| >18 ≤ 40 | 4320 | 81.39 ± 28.89 | 80.56-82.28 | 0.1111 | |
| >40 ≤ 60 | 10929 | 80.52 ± 29.85 | 79.96-81.08 | | |
| >60 | 7582 | 81.30 ± 29.15 | 80.65-81.92 | | |

*Two-sample t-test.

**One-way ANOVA.

high binding affinity for non-specific receptors such as muscarinic (mACh), histaminic (HA-1) and alpha-1 adrenoceptors, which are responsible for side effects such as dry mouth, constipation and sedation [57]. Similarly, paroxetine demonstrates an affinity for mACh receptors similar to that of imipramine and has marked anticholinergic adverse effects [57]. Our data infer that the reduced

compliance with these antidepressants may have a biological basis, in particular an apparent increase in cholinergic sensitivity in the HIV/AIDS population. Indeed, cholinomimetic antibodies of the immunoglobulin (IgA) class are present in HIV/AIDS patients [58,59], suggesting that these patients may indeed be hypersensitive to drugs with activity on the cholinergic system [60]. Importantly,

Table 3 The influence of AD class on AD compliance in MDD patients with/without HIV/AIDS (dispensed items)

| MDD patients with HIV/AIDS (dispensed items) | | | | | |
|---|------|----------------|------------------|---------|------------|
| Antidepressant class | n | Acceptable (%) | Unacceptable (%) | P*** | Cramer's V |
| SSRIs | 207 | 44.93 | 55.07 | <0.0001 | 0.3450 |
| Diverse antidepressants | 148 | 52.70 | 37.30 | | |
| SNRIs | 70 | 58.57 | 41.43 | | |
| TCA | 41 | 26.83 | 73.17 | | |
| MDD patients without HIV/AIDS (dispensed items) | | | | | |
| Antidepressant class | n | Acceptable (%) | Unacceptable (%) | P*** | Cramer's V |
| SSRIs | 9435 | 51.64 | 48.36 | 0.0001 | 0.0942 |
| SNRIs | 5777 | 56.36 | 43.64 | | |
| Diverse antidepressants | 5460 | 48.24 | 51.76 | | |
| TCA | 2159 | 43.72 | 56.28 | | |

***Chi-square.

Table 4 Percentage of the top 10 most frequently dispensed antidepressants with acceptable vs. non-acceptable compliance in MDD patients with/without HIV/AIDS (dispensed items)

| | HIV + MDD | | | | MDD | | | | P*** | Cramer's V |
|---------------|-----------|------------|----|--------------|------|------------|------|--------------|--------|------------|
| | n | Acceptable | n | Unacceptable | n | Acceptable | n | Unacceptable | | |
| Mirtazapine | 40 | 51.28 | 38 | 48.72 | 848 | 46.41 | 979 | 53.59 | 0.3987 | 0.0193 |
| Escitalopram | 34 | 44.16 | 43 | 55.84 | 1781 | 52.74 | 1596 | 47.26 | 0.1359 | -0.0254 |
| Venlafaxine | 30 | 75.00 | 10 | 25.00 | 1972 | 56.93 | 1492 | 43.07 | 0.0217 | 0.0388 |
| Citalopram | 25 | 51.02 | 18 | 48.92 | 1002 | 51.46 | 945 | 48.54 | 0.9511 | -0.0014 |
| Fluoxetine | 19 | 41.30 | 25 | 58.70 | 724 | 47.95 | 786 | 52.05 | 0.3743 | -0.0225 |
| Bupropion | 18 | 78.26 | 5 | 21.74 | 714 | 50.46 | 701 | 49.54 | 0.0082 | 0.0698 |
| Duloxetine | 11 | 36.67 | 19 | 63.33 | 1284 | 55.51 | 1029 | 44.49 | 0.0391 | -0.0426 |
| Amitriptyline | 8 | 23.53 | 26 | 76.47 | 548 | 39.34 | 845 | 60.66 | 0.0618 | -0.0494 |
| Trazodone | 6 | 27.27 | 16 | 72.73 | 596 | 50.72 | 579 | 49.28 | 0.0293 | -0.0630 |
| Paroxetine | 6 | 28.57 | 15 | 71.43 | 582 | 54.19 | 492 | 45.81 | 0.0197 | -0.0705 |

***Chi-square.

TCA and paroxetine non-compliance with TCAs and paroxetine can evoke a cholinergic overdrive [57,61], which may augment the hyper-cholinergic state present in HIV/AIDS patients, leading to a greater adverse experience and subsequent non-compliance. Furthermore, since MDD is associated with increased cholinergic drive [62,63], it is clear that poor compliance in this population may worsen the mood disorder. For the successful treatment of both MDD and HIV/AIDS, compliance with treatment is therefore of utmost importance. Poor compliance with treatment can lead to a drastic decline in quality of life, increased social impairment, low occupational functioning and heightened social isolation, increased suicide ideation and ultimately to suicide [21,55,56,64-66].

Another interesting result is that the class of antidepressant plays a significant role in predicting compliance when treating patients with MDD and HIV/AIDS. The tricyclic antidepressant (TCA) class is associated with the weakest compliance ratio (26.83%), whereas the SNRIs present with the highest compliance ratio (58.57%). According to our results, SSRIs represent the class of ADs with the highest prescription frequency (44.42%, N = 466). Both escitalopram and citalopram are considered as first-line treatment for treating MDD in HIV/AIDS patients because of a limited effect on the cytochrome P450 system, thereby reducing drug-drug interactions with ARV treatment [67-69]. Moreover, a number of studies have proposed that MDD associated with HIV/AIDS should be treated with fluoxetine and paroxetine [70-72]. However, their use is limited by drug-drug interaction with protease inhibitors (PIs) due to their potent inhibition of CYP2D6, thereby increasing levels of PIs and associated toxicity [57,67,73]. However, in this study, the SSRIs as a group displayed a relatively weak compliance ratio (44.93%; $p < 0.0001$; Table 3) compared to the SNRIs, which displayed a significantly

higher compliance ratio (58.57%; $p < 0.0001$; Table 3). These findings indicate that the SSRIs have more adverse effects in this population than originally predicted, possibly related to their anticholinergic (paroxetine), antihistaminic (citalopram) and general serotonergic properties [57]. In fact, considering the latter point, neuropsychiatric symptoms may be evident in HIV-treated patients, especially those on efavirenz, and respond to treatment with the antiserotonergic agent, cyproheptadine, suggesting increased serotonergic activity/sensitivity in these patients [74]. Furthermore, inappropriate AD discontinuation and non-compliance are also associated with increased serotonergic activity [75] that may amplify the adverse experience associated with HIV/AIDS-related neurotransmitter dysregulation. What is more important considering our data is that SNRIs are currently considered a second-line treatment for MDD in HIV/AIDS patients [68,69], while our data (preliminary as they may be) suggest a possible re-evaluation of these guidelines. Indeed, further studies in this regard are warranted.

Only 127 patients (i.e. 0.24% of the total population in the database) met the criteria of co-morbid MDD and HIV/AIDS. Currently, there exists great controversy in the literature regarding the co-morbidity between depression and HIV/AIDS, with a number of articles placing the co-morbidity rate between 0 and 54% [4,6,7,55]. This finding again highlights the complicated co-morbidity between MDD and HIV/AIDS, the different approaches, methodology and reporting of results used in several studies with regard to the selection of a study population, the demographics and the behaviour of the defined population. Furthermore, the low prevalence rate found in this study might suggest a severe under diagnoses of MDD in HIV/AIDS patients and that MDD symptoms might be mistaken rather as symptoms of HIV/AIDS [76,77].

We also observed that female patients ($n = 90$) were more than double that of male patients ($n = 37$). Female patients diagnosed with both MDD and HIV/AIDS displayed significantly lower compliance with AD treatment when compared to male patients of the same group, and displayed almost double the prevalence of both MDD and HIV/AIDS than male counterparts. Some of the reasons given for the poor compliance rate to AD treatment in women are the risk of adverse effects (e.g. weight gain and sexual dysfunction), low socio-economic status, younger age, drug abuse and divorce marital status [78]. Similar to our findings, two separate studies found that almost double the number of women with MDD and HIV/AIDS are likely to die from an HIV/AIDS-related cause when compared to HIV-positive women without depression [79], while females have a significantly higher frequency of MDD and HIV/AIDS vs. males with a combined diagnosis [80,81]. Some authors have suggested that lower levels of testosterone might play a role in the large prevalence of MDD in females when compared to men [82]. In support of this, males with hypogonadism have a significantly higher prevalence of MDD and anxiety disorders when compared to healthy males [83,84]. Moreover, testosterone-replacement therapy in hypogonadal men significantly improves these symptoms [83,85,86]. Importantly, studies have found benefits in administering testosterone to depressed women [87,88], thereby supporting the role of testosterone as a causal factor for MDD in women. However, a study in depressed HIV/AIDS patients has shown that testosterone displayed no significant effect on mood [56].

Examining individual ADs dispensed to patients who are suffering from both MDD and HIV/AIDS, mirtazapine was the most frequently dispensed AD ($n = 78$), followed closely by escitalopram ($n = 77$). Although mirtazapine is not a first-line AD for the treatment of MDD, its side effect profile seems to play a valuable role in late-stage HIV/AIDS patients with co-morbid MDD, with side effects such as constipation, weight gain and sedation helping to improve diarrhoea, diminished appetite and insomnia, respectively [67].

The strength of the current study is that only patients diagnosed with MDD by a psychiatrist were included in the study population. This article has shed light on the current practice of AD treatment in HIV/AIDS patients with medical aids. The most important findings include that HIV/AIDS with co-morbid MDD have reduced compliance with ADs, especially those exhibiting multi-receptor pharmacology, most notably TCAs and certain SSRIs such as paroxetine. Secondly, MDD patients with co-morbid HIV/AIDS display significantly higher compliance with venlafaxine, contrary to current clinical practice where guidelines suggest SSRIs and SNRIs as first and second line treatment options, respectively. Further clarification with

respect to duloxetine is needed as it performed poorly in our analysis. Lastly, female patients are more prone to develop MDD and are notably less compliant with AD treatment.

The authors recognise a few limitations in the current study. Firstly, the numbers of patients with both MDD and HIV/AIDS were relatively small and not a full representation of the national population. Several factors contributed to the small sample size, such as the use of very strict inclusion criteria, only a portion of all the medical aids is listed with the PBM utilised in this study, depressed patients using psychological treatment for MDD were not taken into account, while the study population consisted of only patients contributing to a private medical aid/insurance. Secondly, the use of a database limits the further extraction of patient information regarding the stage of HIV/AIDS in which the patient is and whether there was an improvement or deterioration in the patient's mood.

Conclusion

This study confirms that AD compliance is lower in depressed HIV/AIDS vs. depressed non-HIV/AIDS patients, correlating with global trends regarding non-compliance of patients with a chronic disease and co-morbid disorders. However, this study indicated that the clinical and practical significance of the differences is low. Moreover, the fact that the prevalence of MDD patients with co-morbid HIV/AIDS was limited in this study again highlights the need for more research in various population groups with a focus on the socio-economic status of the patients. Although our data are preliminary and warrant further study, it does reveal evidence that venlafaxine may be a valid choice of antidepressant in HIV/AIDS patients suffering from MDD and should be considered as first-line antidepressant in such cases. Moreover, our data suggest that multi-target antidepressants such as TCAs and select SSRIs may increase the risk of non-compliance in this population. Lastly, when prescribing ADs to females, special consideration should be given to counselling them on both their illness and treatment so that they clearly understand the benefits of compliance to treatment.

Methods

We conducted a prospective, cohort study analysing nationally representative medicine claims data submitted to a privately-owned South African Pharmaceutical Benefit Management (PBM) company. The data represent a third of South African patients with private medical aids. No distinction was made between races. Two groups were distinguished in the database over a six-year study period (1 January 2006 to 31 December 2011), namely patients with only MDD ($n = 12\ 270$) and MDD patients with HIV/AIDS ($n = 127$).

We queried data for patient demographics (gender and date of birth) and pertinent prescription information (such as drug trade name, days supplied, dispensing date, quantity of medicine prescribed, and ICD-10 code per claim). The following automated validation processes were applied by the PBM that ascertained the quality of data: data integrity validation, eligibility management, medicine utilisation and clinical management, pricing and formulary management. There were no missing data fields in the datasets. The variables 'birth date' and 'dispensing date' were used to calculate the age of patients on the date of treatment and the number of days between refills.

A study population was selected according the following inclusion criteria: patients older than 18 years, MDD should be diagnosed by a psychiatrist supported by an appropriate ICD-10 code. The ICD-10 codes are based on the International Classification of Diseases, 10th edition published by the World Health Organization [89]. In this study, the ICD-10 codes F32 (Depressive episode) and F33 (Recurrent depressive disorder) were used to identify patients with MDD as diagnosed by a psychiatrist, as well as B20-B24 (Human immunodeficiency virus [HIV/AIDS] disease), and all patients on cARV treatment. Thereby it was ensured that data were excluded where ADs may have been used for other indications, such as amitriptyline for the treatment of chronic pain.

For the purpose of this study, the medicine possession ratio (MPR) was used to determine AD compliance of patients. The MPR is a well-established method to calculate drug compliance in pharmacoepidemiological studies, including chronic diseases such as MDD [90], hypertension [91], and schizophrenia [92]. However, it is important to note that the compliance value obtained from the MPR only gives an indication of the possession of medicine by the patient, and that appropriate consumption of medicine is assumed to ensue from possession. The usage of medicine claims data to determine MPR calculations is valuable in that it is acceptably accurate, convenient, objective, non-invasive and relatively inexpensive to obtain when a large study population is needed. It is therefore suitable for the calculation of MPR as an indication of patient compliance with medication therapy [93,94]. However, the MPR has a number of limitations. Firstly, MPR can only be calculated if a patient has filled more than two prescriptions. Secondly, MPR can only assess whether the medication was used consistently. Lastly, the MPR cannot measure whether a patient was compliant with the instructions given by the medical practitioner [95].

The MPR is defined as the number of days for which medication is supplied within the refill interval (medicine

treatment period) divided by the number of days in the refill interval [96,97].

$$\text{MPR} = \frac{\text{Number of days in refill interval}}{\text{Sum the days of supplied medication}} \times 100$$

The MPR is considered acceptable if the calculated value is $\geq 80\%$, but $\leq 110\%$. An MPR of less than 80% indicates the presence of refill gaps so that possession is considered unacceptably low (undersupply), whereas an MPR greater than 110% is considered unacceptably high (oversupply). Conversely, a patient was considered compliant (acceptable group) with his/her AD treatment if the MPR was $\geq 80\%$ and $\leq 110\%$, and AD treatment period was longer than 120 days. Therefore, distinguishing between compliant MPR group and patients out of this range will be considered as the non-compliant MPR group or unacceptable group.

Data management and analysis were performed in SAS Version 9.1.3 (SAS Institute, Cary, NC) [98]. All statistical significance was considered with a probability of $p < 0.05$. The practical significance of results was computed when the p -value was statistically significant ($p \leq 0.05$).

Variables (age groups, gender and active ingredients) were expressed using descriptive statistics such as frequencies (n), percentages (%), means, standard deviations and 95% confidence intervals (CI). The patient's age was determined at time of first dispensing and divided into three groups: > 18 to ≤ 40 years; > 40 to ≤ 60 years; > 60 years.

The dataset is very large, therefore all the statistical methods used in this study rely on the Central Limit Theorem, which states that the average of a large number of independent random variables is approximately normally distributed around the true population mean [99]. Therefore, the two-sample t-test allowed us to compare the mean MPR of male and female patients. The one-way ANOVA was used to test differences between the mean MPR of different age groups and AD classes. It was operationalised with the general linear model (GLM) procedure of the SAS Version 9.1.3. If a difference was indicated, a Tukey's multiple comparison test was performed to determine which groups differ significantly from each other. Cohen's d was used to evaluate effect size between means (with $d \geq 0.8$ defined as a large effect with practical significance). The Chi-square test (χ^2) was used to determine whether an association exists between proportions of two or more groups (compliance vs. active ingredients). The Cramer's V statistic was used to test the practical significance of this association (with Cramer's $V \geq 0.5$ defined as practical significance).

This study was approved by the Research Ethics Committee of the North-West University (NWU-0046-08-550) and the Board of Directors of the South African Pharmaceutical Benefit Management (PBM). Data were analysed anonymously.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

FN Slabbert has been involved in the design of the study, drafted the manuscript, and carried out the analyses and interpretation of data. MS Lubbe made a substantial contribution in the design of the study and developed the methodology. BH Harvey and CB Brink revised the manuscript and have given extensive intellectual inputs. All authors read and approved the final manuscript.

Authors' information

FN Slabbert: BPharm, MSc (Pharmacology). PhD student and junior research scientist, School of Pharmacy, North-West University (Potchefstroom Campus). MS Lubbe: BPharm, MPharm, PhD (Pharmacy Practice). Leader: Medicine Usage in South Africa, professor at the School of Pharmacy, North-West University (Potchefstroom Campus). BH Harvey: BPharm, BSc (Hons), MSc (Pharmacology), PhD (Pharmacology). Center of Excellence for Pharmaceutical Sciences, Co-director – MRC unit on Anxiety and Stress Disorders. Professor of Pharmacology, School of Pharmacy, North-West University (Potchefstroom Campus). CB Brink: BPharm, MSc (Pharmacology), PhD (Pharmacology). Secretary-general: WCP2014. President: SASBCP. Chair: IUPHAR-Ed. Professor of Pharmacology, School of Pharmacy, North-West University (Potchefstroom Campus).

Acknowledgements

We thank Dr Suria Ellis from Statistical Consultation Services and Mrs Marike Cockeran from Medicine Usage in South Africa, North-West University (Potchefstroom Campus) for statistical support, and Anne-Marie Bekker for administrative support regarding the database. The authors acknowledge the North-West University, National Research Foundation and the South African Medical Research Council for financial support.

Author details

¹Medicines Usage Group (MUSA), North-West University, Potchefstroom 2520, South Africa. ²Division of Pharmacology, North-West University, Potchefstroom 2520, South Africa. ³Centre of Excellence for Pharmaceutical Sciences, School of Pharmacy, North-West University, Potchefstroom 2520, South Africa.

Received: 2 July 2014 Accepted: 2 March 2015

Published online: 14 April 2015

References

- UNAIDS. UNAIDS World AIDS Day Report 2012; UNAIDS Fact Sheet 2012; Kaiser Family Foundation. In Book UNAIDS World AIDS Day Report 2012; UNAIDS Fact Sheet 2012; Kaiser Family Foundation. (Editor ed. eds.). City; 2012 [http://www.unaids.org/sites/default/files/en/media/unaids/contentassets/documents/epidemiology/2012/gr2012/20121120_FactSheet_Global_en.pdf]
- Centers for Disease Control and Prevention. HIV surveillance report: Diagnoses of HIV infection and AIDS in the United States and dependent areas, 2009. [http://www.cdc.gov/hiv]
- World Health Organisation. Data on the size of the HIV/AIDS epidemic: Number of people (all ages) living with HIV by country [http://apps.who.int/gho/data/view.main.22100]
- Kinyanda E, Hoskins S, Nakku J, Nawaz S, Patel V. Prevalence and risk factors of major depressive disorder in HIV/AIDS as seen in semi-urban Entebbe district, Uganda. *BMC Psychiatry*. 2011, 11.
- Olley BO, Seedat S, Stein DJ. Persistence of psychiatric disorders in a cohort of HIV/AIDS patients in South Africa: a 6-month follow-up study. *J Psychosom Res*. 2006;61:479–84.
- Cysique LA, Deutsch R, Atkinson JH, Young C, Marcotte TD, Dawson L, et al. Incident major depression does not affect neuropsychological functioning in HIV-infected men. *J Int Neuropsychol Soc*. 2006;13:1–11.
- Gibbie T, Hay M, Hutchison CW, Mich A. Depression, social support and adherence to highly active antiretroviral therapy in people living with HIV/AIDS. *Sex Health*. 2007;4:227–32.
- Ali S, Stone MA, Peters JL, Davies MJ, Khunti K. The prevalence of co-morbid depression in adults with Type 2 diabetes: a systematic review and meta-analysis. *Diabet Med*. 2006;23:1165–73.
- Schleifer SJ, Macari-Hinson MM. The nature and course of depression following myocardial infarction. *Arch Intern Med*. 1989;149:1785–9.
- Spijkerman T, de Jonge P, van den Brink RH, Jansen JH, May JF, Crijns HJ, et al. Depression following myocardial infarction: first-ever versus ongoing and recurrent episodes. *Gen Hosp Psychiatry*. 2005;27:411–7.
- Von Korff M, Scott KM, Gureje O. Global perspectives on mental-physical comorbidity in the WHO world mental health surveys. Cambridge, UK: Cambridge University Press; 2009.
- Springer SA, Dushaj A, Azar MM. The impact of DSM-IV mental disorders on adherence to combination antiretroviral therapy among adult persons living with HIV/AIDS: a systematic review. *AIDS Behav*. 2012;16:2119–43.
- Katon WJ. Clinical and health services relationships between major depression, depressive symptoms, and general medical illness. *Biol Psychiatry*. 2003;54:216–26.
- DiMatteo MR, Lepper HS, Croghan TW. Depression is a risk factor for noncompliance with medical treatment - Meta-analysis of the effects of anxiety and depression on patient adherence. *Arch Intern Med*. 2000;160:2101–7.
- Plummer ML, Watson-Jones D, Lees S, Baisley K, Matari S, Chagalucha J, et al. A qualitative study of participant adherence in a randomized controlled trial of herpes suppressive therapy for HIV prevention in Tanzania. *AIDS care*. 2010;22:499–508.
- Singh N, Berman SM, Swindells S, Justis JC, Mohr JA, Squier C, et al. Adherence of human immunodeficiency virus-infected patients to antiretroviral therapy. *Clin Infect Dis*. 1999;29:824–30.
- Ellis RJ, Rosario D, Clifford DB, McArthur JC, Simpson D, Alexander T, et al. Continued high prevalence and adverse clinical impact of human immunodeficiency virus-associated sensory neuropathy in the era of combination antiretroviral therapy: the CHARTER Study. *Arch Neurol*. 2010;67:552–8.
- Orlando M, Burnam MA, Beckman R, Morton SC, London AS, Bing EG, et al. Re-estimating the prevalence of psychiatric disorders in a nationally representative sample of persons receiving care for HIV: results from the HIV Cost and Services Utilization Study. *Int J Methods Psychiatr Res*. 2002;11:75–82.
- Lima VD, Geller J, Rangsberg DR, Patterson TL, Daniel M, Kerr T, et al. The effect of adherence on the association between depressive symptoms and mortality among HIV-infected individuals first initiating HAART. *AIDS*. 2007;21:1175–83.
- Sin NL, DiMatteo MR. Depression treatment enhances adherence to antiretroviral therapy: a meta-analysis. *Ann Behav Med*. 2014;47:259–69.
- Alciati A, Starace F, Scaramelli B, Campaniello M, Adriani B, Mellado C, et al. Has there been a decrease in the prevalence of mood disorders in HIV-seropositive individuals since the introduction of combination therapy? *Eur Psychiatry*. 2001;16:491–6.
- Ickovics JR, Hamburger ME, Vlahov D, Schoenbaum EE, Schuman P, Boland RJ, et al. Mortality, CD4 cell count decline, and depressive symptoms among HIV-seropositive women - Longitudinal analysis from the HIV epidemiology research study. *JAMA*. 2001;285:1466–74.
- Lyketsos CG, Hoover DR, Guccione M, Dew MA, Wesch JE, Bing EG, et al. Changes in depressive symptoms as AIDS develops. The Multicenter AIDS Cohort Study. *Am J Psychiatry*. 1996;153(11):1430–7.
- Lyketsos CG, Hoover DR, Guccione M. Depression and survival among HIV-infected persons. *JAMA*. 1996;275:35–6.
- Lopes M, Olsson M, Rabkin J, Hasin DS, Alegria AA, Lin KH, et al. Gender, HIV status, and psychiatric disorders: Results from the National Epidemiologic Survey on Alcohol and Related Conditions. *J Clin Psychiatry*. 2012;73:384–91.
- Schuster R, Bornova M, Hunt E. The influence of depression on the progression of HIV: direct and indirect effects. *Behav Modif*. 2012;36:123–45.
- Harvey BH, Meyer CL, Gallichio VS, Manji HK. Lithium salts in AIDS and AIDS-related dementia. *Psychopharmacol Bull*. 2002;36:5–26.
- Leonard BE. The concept of depression as a dysfunction of the immune system. *Curr Immunol Rev*. 2010;6:205–12.
- Weinstock M. Intrauterine factors as determinants of depressive disorder. *Isr J Psychiatr Rel*. 2010;47:36–45.
- Loftis JM, Huckans M, Morasco BJ. Neuroimmune mechanisms of cytokine-induced depression: current theories and novel treatment strategies. *Neurobiol Dis*. 2010;37:519–33.
- Khairova RA, Machado-Vieira R, Du J, Manji HK. A potential role for pro-inflammatory cytokines in regulating synaptic plasticity in major depressive disorder. *Int J Neuropsychopharmacol*. 2009;12:561–78.
- Krishnadas R, Cavanagh J. Depression: an inflammatory illness? *J Neurol Neurosurg Psychiatry*. 2012;83:495–502.

33. Capuron L, Ravaud A, Dantzer R. Early Depressive Symptoms in Cancer Patients Receiving Interleukin 2 and/or Interferon Alfa-2b Therapy. *J Clin Oncol*. 2000;18:2143–51.
34. Capuron L, Ravaud A, Miller AH, Dantzer R. Baseline mood and psychosocial characteristics of patients developing depressive symptoms during interleukin-2 and/or interferon-alpha cancer therapy. *Brain Behav Immun*. 2004;18:205–13.
35. Sotelo JL, Musselman D, Nemeroff C. The biology of depression in cancer and the relationship between depression and cancer progression. *Int Rev Psychiatry*. 2014;26:16–30.
36. Del Guerra FB, Fonseca JL, Figueiredo VM, Ziff EB, Konkiewitz EC. Human immunodeficiency virus-associated depression: contributions of immunoinflammatory, monoaminergic, neurodegenerative, and neurotrophic pathways. *J Neurovirol*. 2013;19:314–27.
37. Crum NF, Riffenburgh RH, Wegner S, Agan BK, Tasker SA, Spooner KM, et al. Comparisons of causes of death and mortality rates among HIV-infected persons: Analysis of the pre-, early, and late HAART (highly active antiretroviral therapy) eras. *J Acquir Imm Def*. 2006;41:194–200.
38. Lima VD, Harrigan R, Bangsberg DR, Hogg RS, Gross R, Yip B, et al. The combined effect of modern highly active antiretroviral therapy regimens and adherence on mortality over time. *J Acquir Immune Defic Syndr*. 2009;50:529–36.
39. Bhaskaran K, Hamouda O, Sannes M, Boufassa F, Johnson AM, Lambert PC, et al. Changes in the risk of death after HIV seroconversion compared with mortality in the general population. *JAMA*. 2008;300:51–9.
40. Bae JW, Guyer W, Grimm K, Altice FL. Medication persistence in the treatment of HIV infection: a review of the literature and implications for future clinical care and research. *AIDS*. 2011;25:279–90.
41. de Bruin M, Hospers HJ, van Breukelen GJ, Kok G, Koevoets WM, Prins JM. Electronic monitoring-based counseling to enhance adherence among HIV-infected patients: a randomized controlled trial. *Health Psychol*. 2010;29:421–8.
42. Angelino AF, Treisman GJ. Issues in co-morbid severe mental illnesses in HIV infected individuals. *Int Rev Psychiatry*. 2008;20:95–101.
43. Treisman GJ, Angelino AF, Hutton HE. Psychiatric issues in the management of patients with HIV infection. *JAMA*. 2001;286:2857–64.
44. Paterson DL, Swindells S, Mohr J, Brester M, Vergis EN, Squier C, et al. Adherence to protease inhibitor therapy and outcomes in patients with HIV infection. *Ann Intern Med*. 2000;133:21–30.
45. Nachega JB, Hislop M, Dowdy DW, Chaisson RE, Regensberg L, Maertens G. Adherence to nonnucleoside reverse transcriptase inhibitor-based HIV therapy and virologic outcomes. *Ann Intern Med*. 2007;146:564–73.
46. Mannheimer S, Friedland G, Matts J, Child C, Chesney M. The consistency of adherence to antiretroviral therapy predicts biologic outcomes for human immunodeficiency virus-infected persons in clinical trials. *Clin Infect Dis*. 2002;34:1115–21.
47. Nieuwkerk PT, Oort FJ. Self-reported adherence to antiretroviral therapy for HIV-1 infection and virologic treatment response: A meta-analysis. *J Acquir Immune Defic Syndr*. 2005;38:445–8.
48. Gonzalez JS, Batchelder AW, Psaros C, Safren SA. Depression and HIV/AIDS treatment nonadherence: a review and meta-analysis. *J Acquir Immune Defic Syndr*. 2011;58:181–7.
49. Marcus EN. The silent epidemic: The health effects of illiteracy. *N Engl J Med*. 2006;355:339–41.
50. Chesney MA. The elusive gold standard. Future perspectives for HIV adherence assessment and intervention. *J Acquir Immune Defic Syndr*. 2006;43 Suppl 1:S149–155.
51. Schneider J, Kaplan SH, Greenfield S, Li W, Wilson IB. Better physician-patient relationships are associated with higher reported adherence to antiretroviral therapy in patients with HIV infection. *J Gen Intern Med*. 2004;19:1096–103.
52. Nierenberg AA. Long-term management of chronic depression. *J Clin Psychiatry*. 2001;62 Suppl 6:17–21.
53. Warner CH, Bobo W, Warner C, Reid S, Rachal J. Antidepressant discontinuation syndrome. *Am Fam Physician*. 2006;74:449–56.
54. Akerblad AC, Bengtsson F, Holgersson M, von Knorring L, Ekselius L. Identification of primary care patients at risk of nonadherence to antidepressant treatment. *Patient Prefer Adherence*. 2008;2:379–86.
55. Chandler G, Himelhoch S, Moore RD. Substance abuse and psychiatric disorders in HIV-positive patients: Epidemiology and impact on antiretroviral therapy. *Drugs*. 2006;66(6):769–89.
56. Rabkin JG. HIV and depression: 2008 review and update. *Curr HIV/AIDS Rep*. 2008;5:163–71.
57. Harvey BH. The neurobiology and pharmacology of depression. A comparative overview of serotonin selective antidepressants. *S Afr Med J*. 1997;87:540–50. 552.
58. Borda E, Sterin-Borda L, Vernava D, Rodriguez M, Perez-Bianco R, De Bracco MME. Parasympathomimetic activity of sera from HIV-infected hemophilic patients with or without diarrhea. *Immunol Infect Dis*. 1993;3:321–6.
59. de Bracco MM, Borda E, Galassi N, Perez-Bianco R, Sterin-Borda L. Autoantibodies in HIV-infected patients that modulate the cholinergic activity of heart and gut tissue. *Autoimmunity*. 1993;14:307–14.
60. Sales ME, Sterin-Borda L, Bracco MME, Rodriguez M, Narbaitz M, Borda E. IgA from HIV+ haemophilic patients triggers intracellular signals coupled to the cholinergic system of the intestine. *Clin Exp Immunol*. 1997;110:189–95.
61. Harvey BH, McEwen BS, Stein DJ. Neurobiology of antidepressant withdrawal: implications for the longitudinal outcome of depression. *Biol Psychiatry*. 2003;54:1105–17.
62. Furey ML, Khanna A, Hoffman EM, Drevets WC. Scopolamine produces larger antidepressant and anti-anxiety effects in women than in men. *Neuropsychopharmacology*. 2010;35:2479–88.
63. Mineur YS, Obayemi A, Wigstrand MB, Fote GM, Calarco CA, Li AM, et al. Cholinergic signaling in the hippocampus regulates social stress resilience and anxiety- and depression-like behavior. *Proc Natl Acad Sci U S A*. 2013;110:3573–8.
64. Carrico AW, Antoni MH, Duran RE, Ironson G, Penedo F, Fletcher MA, et al. Reductions in depressed mood and denial coping during cognitive behavioral stress management with HIV-positive gay men treated with HAART. *Ann Behav Med*. 2006;31:155–64.
65. Cooperman NA, Simoni JM. Suicidal ideation and attempted suicide among women living with HIV/AIDS. *J Behav Med*. 2005;28:149–56.
66. Haller DL, Miles DR. Suicidal ideation among psychiatric patients with HIV: Psychiatric morbidity and quality of life. *AIDS Behav*. 2003;7:101–8.
67. Thompson A, Silverman B, Dzung L, Treisman G. Psychotropic medications and HIV. *Clin Infect Dis*. 2006;42:1305–10.
68. Freudenreich O, Goforth HW, Cozza KL, Mimiaga MJ, Safren SA, Bachmann G, et al. Psychiatric treatment of persons with HIV/AIDS: An HIV-psychiatry consensus survey of current practices. *Psychosomatics*. 2010;51:480–8.
69. Relf MV, Eibach S, Okine KN, Ward T. Evidence-based clinical practice guidelines for managing depression in persons living with HIV. *J Assoc Nurses AIDS Care*. 2013;24:S15–28.
70. Caballero J, Nahata MC. Use of selective serotonin-reuptake inhibitors in the treatment of depression in adults with HIV. *Ann Pharmacother*. 2005;39:141–5.
71. Elliott AJ, Uldall KK, Bergam K, Russo J, Claypoole K, Roy-Byrne PP. Randomized, placebo-controlled trial of paroxetine versus imipramine in depressed HIV-positive outpatients. *Am J Psychiatry*. 1998;155:367–72.
72. Treisman G, Fishman M, Lyketsos C, McHugh PR. Evaluation and treatment of psychiatric disorders associated with HIV infection. *Res Publ Assoc Res Nerv Ment Dis*. 1994;72:239–50.
73. Yanofski J, Croarkin P. Choosing antidepressants for HIV and AIDS patients: Insights on safety and side effects. *Psychiatry (Edgmont)*. 2008;5:61–6.
74. Dabaghzadeh F, Ghaeli P, Khalili H, Alimadadi A, Jafari S, Akhondzadeh S, et al. Cyproheptadine for prevention of neuropsychiatric adverse effects of efavirenz: a randomized clinical trial. *AIDS Patient Care STDS*. 2013;27:146–54.
75. Harvey BH, Slabbert FN. New insights on the antidepressant discontinuation syndrome. *Hum Psychopharmacol*. 2014, in press.
76. Asch SM, Kilbourne AM, Gifford AL, Burnam MA, Turner B, Shapiro MF, et al. Underdiagnosis of depression in HIV: Who are we missing? *J Gen Intern Med*. 2003;18:450–60.
77. Pyne JM, Asch SM, Lincourt K, Kilbourne AM, Bowman C, Atkinson H, et al. Quality indicators for depression care in HIV patients. *AIDS Care*. 2008;20:1075–83.
78. Demyttenaere K. Compliance during treatment with antidepressants. *J Affect Disord*. 1997;43:27–39.
79. Cook JA, Grey D, Burke J, Cohen MH, Gurtman AC, Richardson JL, et al. Depressive symptoms and AIDS-related mortality among a multisite cohort of HIV-positive women. *Am J Public Health*. 2004;94:1133–40.
80. McKnight-Eily LR, Presley-Cantrell L, Elam-Evans LD, Chapman DP, Kaslow NJ, Pery GS. Prevalence and correlates of current depressive symptomatology and lifetime diagnosis of depression in black women. *Womens Health Issues*. 2009;19:243–52.

81. Weissman MM, Bland RC, Canino GJ, Faravelli C, Greenwald S, Hwu HG, et al. Cross-national epidemiology of major depression and bipolar disorder. *JAMA*. 1996;276:293–9.
82. McHenry J, Carrier N, Hull E, Kabbaj M. Sex differences in anxiety and depression: role of testosterone. *Front Neuroendocrinol*. 2014;35:42–57.
83. Zarrouf FA, Artz S, Griffith J, Sirbu C, Kommor M. Testosterone and depression: systematic review and meta-analysis. *J Psychiatr Pract*. 2009;15:289–305.
84. DiBlasio CJ, Hammett J, Malcolm JB, Judge BA, Womack JH, Kincade MC, et al. Prevalence and predictive factors for the development of de novo psychiatric illness in patients receiving androgen deprivation therapy for prostate cancer. *Can J Urol*. 2008;15:4249–56. discussion 4256.
85. Pope HG, Cohane GH, Kanayama G, Siegel AJ, Hudson JL. Testosterone gel supplementation for men with refractory depression: A randomized, placebo-controlled trial. *Am J Psychiatr*. 2003;160:105–11.
86. Wang C, Alexander G, Berman N, Salehian B, Davidson T, McDonald V, et al. Testosterone replacement therapy improves mood in hypogonadal men: A clinical research center study. *J Clin Endocrinol Metab*. 1996;81:3578–83.
87. Miller KK, Perlis RH, Papakostas GI, Mischoulon D, Losifescu DV, Brick DJ, et al. Low-dose transdermal testosterone augmentation therapy improves depression severity in women. *CNS Spectr*. 2009;14:688–94.
88. Shifren JL, Braunstein GD, Simon JA, Casson PR, Buster JE, Redmond GP, et al. Transdermal testosterone treatment in women with impaired sexual function after oophorectomy. *N Engl J Med*. 2000;343:682–8.
89. World Health Organization: ICD-10 Version: 2010. [<http://www.who.int/classifications/icd/icdonlineversions/en/>]
90. Serna MC, Cruz I, Real J, Gasco E, Galvan L. Duration and adherence of antidepressant treatment (2003 to 2007) based on prescription database. *Eur Psychiatry*. 2010;25:206–13.
91. Cramer JA, Benedict A, Muszbek N, Keskinaslan A, Khan ZM. The significance of compliance and persistence in the treatment of diabetes, hypertension and dyslipidaemia: A review. *Int J Clin Pract*. 2008;62:76–87.
92. Weiden PJ, Kozma C, Grogg A, Locklear J. Partial compliance and risk of rehospitalization among California Medicaid patients with schizophrenia. *Psychiatr Serv*. 2004;55:886–91.
93. Medic G, Higashi K, Littlewood KJ, Diez T, Granstrom O, Kahn RS. Dosing frequency and adherence in chronic psychiatric disease: systematic review and meta-analysis. *Neuropsychiatr Dis Treat*. 2013;9:119–31.
94. Prukkanone B, Vos T, Burgess P, Chaiyakunapruk N, Bertram M. Adherence to antidepressant therapy for major depressive patients in a psychiatric hospital in Thailand. *BMC Psychiatry*. 2010;10:64.
95. Zhao B, Wong EC, Palaniappan L. Estimating patient adherence to medication with electronic health records data and pharmacy claims combined. *SAS Global Forum 2013*. [<http://support.sas.com/resources/papers/proceedings13/167-2013.pdf>]
96. Steiner JF, Prochazka AV. The assessment of refill compliance using pharmacy records: methods, validity, and applications. *J Clin Epidemiol*. 1997;50:105–16.
97. Steyn, HS. Manual for the determination of effect size indices and practical significance. Potchefstroom, North-West University. <http://www.nwu.ac.za/content/stats-effect-size>. Date of access: 25 Sep. 2012.
98. SAS Institute Inc., 2002-2010. SAS for Windows 9.3 Cary, NC.
99. Lumley T, Diehr P, Emerson S, Chen L. The importance of the normality assumption in large public health data sets. *Annu Rev Public Health*. 2002;23:151–69.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

