

CHAPTER SIX

References

6.1 GENERAL REFERENCES

- Abdalla J.A., Casley W.L., Hudson A.J., Murphy E.G., Cousin H.K., Armstrong H.A. and Ebers G.C. Linkage analysis of candidate loci in autosomal dominant myotonia congenital. *Neurology*, **42**, 1561-1564, 1992.
- ABI PRISM[®] BigDye[™] Terminator version 3.0 Ready Reaction Cycle Sequencing kit, protocol. Applied Biosystems Corporation, Foster City, CA, USA, 2001.
- Adams B., Smith A.T., Bailey S., McEwan A.G. and Bray R.C. Reactions of dimethylsulfoxide reductase from *Rhodobacter capsulatus* with dimethyl sulfide and with dimethyl sulfoxide: Complexities revealed by conventional and stopped-flow spectrophotometry. *Biochemistry*, **38**, 8501-8511, 1999.
- Adeokun A.M., West S.P., Ellis F.R., Halsall P.J., Hopkins P.M., Foughmand A.M., Iles D.E., Robinson R.L., Stewart A.D. and Curran J.L. The G1021A substitution in the RYR1 gene does not cosegregate with malignant hyperthermia susceptibility in a British pedigree. *Am. J. Hum. Genet.*, **60**, 833-841, 1997.
- Adeola O., Ball R.O., House J.D. and O'Brien P.J. Regional brain neurotransmitter concentrations in stress susceptible pigs. *J. Anim. Sci.*, **71**, 968-974, 1993.
- Adnet P.J., Krivosic-Horber R.M., Adamantidis M.M., Haudecoeur G., Adnet-Bonte C.A., Saulnier F. and Dupuis B.A. The association between the neuroleptic malignant syndrome and malignant hyperthermia. *Acta. Anaesthesiol. Scand.*, **33**, 676-680, 1989.
- Adnet P.J., Krivosic-Horber R.M., Adamantidis M.M., Haudecoeur G., Reyford G.H. and Dupuis B.A. Clinical concentrations of verapamil affect the *in vitro* diagnosis of susceptibility to malignant hyperpyrexia. *Br. J. Anaesth.*, **64**, 64-66, 1990a.
- Adnet P.J., Krivosic-Horber R.M., Haudecoeur G., Adamantidis M.M., Reyford G.H., Imbenotte M. and Cordonnier C. Use of the calcium agonist bay K 8644 for *in vitro* diagnosis of susceptibility to malignant hyperthermia. *Br. J. Anaesth.*, **65**, 791-795, 1990b.
- Adnet P.J., Bromberg N.L., Haudecoeur G., Krivosic I., Adamantidis M.M., Reyford H., Bello N. and Krivosic-Horber R.M. Fibre-type caffeine sensitivities in skinned muscle fibres from humans susceptible to malignant hyperthermia. *Anesthesiology*, **78**, 168-177, 1993.
- Adnet P.J., Lestavel P. and Krivosic-Horber R. Neuroleptic malignant syndrome. *Br. J. Anaesth.*, **85**, 129-135, 2000.
- Aleman M., Riehl J., Aldridge B.M., Lecouteur R.A., Stott J.L. and Pessah I.N. Association of a mutation in the ryanodine receptor 1 gene with equine malignant hyperthermia. *Muscle Nerve*, **30**, 356-365, 2004.
- Ali S.Z., Taguchi A. and Rosenberg H. Malignant hyperthermia. *Best Pract. Res. Clin. Anaesthesiol.*, **17**, 519-533, 2003.
- Allen G.C., Cattran C.B., Peterson R.G. and Lalande M. Plasma levels of dantrolene following oral administration in malignant hyperthermia-susceptible patients. *Anesthesiology*, **69**, 900-904, 1988.
- Allen G.C. and Rosenberg H. Malignant hyperthermia susceptibility in adult patients with masseter muscle rigidity. *Can. J. Anaesth.*, **37**, 31-35, 1990.
- Allen G.C. Malignant hyperthermia susceptibility. *Anaesthesiol. Clin. North America*, **12**, 513-535, 1994.
- Allen G.C. and Brubaker C.L. Human malignant hyperthermia associated with desflurane anaesthesia. *Anesth. Analg.*, **86**, 1328-1331, 1998.
- Allen G.C., Larach M.G., Kunselman A.R. and the North American Malignant Hyperthermia registry of MHAUS. The sensitivity and specificity of the caffeine-halothane contracture test. *Anesthesiology*, **88**, 579-588, 1998.
- Alphay L. DNA sequencing from experimental methods to bioinformatics, BIOS Scientific Publishers, Ltd., Oxford, UK, 1997.
- Altschul S.F., Madden T.L., Schäffer A.A., Zhang J., Zhang Z., Miller W. and Lipman D.J. Gapped BLAST and PSI-BLAST: A new generation of protein database search programs. *Nucleic Acids Res.*, **25**, 3389-3402, 1997.

- Anderson T.E., Drummond D.S., Breed A.L. and Taylor C.A. Malignant hyperthermia in myelomeningocele: A previously unreported association. *J. Pediatr. Orthop.*, **1**, 401-403, 1981.
- Anetseder M., Hager M., Müller C.R. and Roewer N. Diagnosis of susceptibility to malignant hyperthermia by use of a metabolic test. *Lancet*, **359**, 1579-1580, 2002.
- Arezi B., Xing W., Sorge J.A. and Hogrefe H.H. Amplification efficiency of thermostable DNA polymerase. *Anal. Biochem.*, **321**, 226-235, 2003.
- Avila G. and Dirksen R.T. Functional effects of central core disease mutations in the cytoplasmic region of the skeletal muscle ryanodine receptor. *J. Gen. Physiol.*, **118**, 277-290, 2001.
- Avila G., O'Connell K.M.S. and Dirksen R.T. The pore region of the skeletal muscle ryanodine receptor is a primary locus for excitation-contraction uncoupling in central core disease. *J. Gen. Physiol.*, **121**, 277-286, 2003.
- Bachand M., Vachon N., Boisvert M., Mayer F.M. and Chartrand D. Clinical reassessment of malignant hyperthermia in Abitibi-Témiscamingue. *Can. J. Anaesth.*, **44**, 696-701, 1997.
- Bäckman E., Lennmarken C., Rutberg H. and Henriksson K.G. Skeletal muscle contraction characteristics *in vivo* in malignant hyperthermia susceptible subjects. *Acta Neurol. Scand.*, **77**, 278-282, 1988.
- Badano J.L., Kim J.C., Hoskins B.E., Lewis R.A., Ansley S.J., Cutler D.J., Castellan C., Beales P.L., Leroux M.R. and Katsanis N. Heterozygous mutations in BBS1, BBS2 and BBS6 have a potential epistatic effect on Bardet-Biedl patients with two mutations at a second BBS locus. *Hum. Mol. Genet.*, **12**, 1651-1659, 2003.
- Ball S.P. and Johnson K.J. The genetics of malignant hyperthermia. *J. Med. Genet.*, **30**, 89-93, 1993.
- Barone V., Massa O., Intravaia E., Bracco A., Di Martino A., Tegazzin V., Cozzolino S. and Sorrentino V. Mutation screening of the RYR1 gene and identification of two novel mutations in Italian malignant hyperthermia families. *J. Med. Genet.*, **36**, 115-118, 1999.
- Beales P.L., Elcioglu N., Woolf A.S., Parker D. and Flinter F.A. New criteria for improved diagnosis of Bardet-Biedl syndrome: Results of a population survey. *J. Med. Genet.*, **36**, 437-446, 1999.
- Beard N.A., Sakowska M.M., Dulhunty A.F. and Laver D.R. Calsequestrin is an inhibitor of skeletal muscle ryanodine receptor calcium release channels. *Biophys. J.*, **82**, 310-20, 2002.
- Beard N.A., Laver D.R. and Dulhunty A.F. Calsequestrin and the calcium release channel of skeletal and cardiac muscle. *Prog. Biophys. Mol. Biol.*, **85**, 33-69, 2004.
- Belmont M.R. Succinylcholine/suxamethonium. *Curr. Opin. Anaesth.*, **8**, 362-366, 1995.
- Benacquista B.L., Sharma M.R., Samsó M., Zorzato F., Treves S. and Wagenknecht T. Amino acid residues 4425-4621 localised on the three-dimensional structure of the skeletal muscle ryanodine receptor. *Biophys. J.*, **78**, 1349-1358, 2000.
- Bernardi G. and Bernardi G. Compositional constraints and genome evolution. *J. Mol. Evol.*, **24**, 1-11, 1986.
- Bernardi G., Mouchiroud D., Gautier C. and Bernardi G. Compositional patterns in vertebrate genomes: Conservation and change in evolution. *J. Mol. Evol.*, **28**, 7-18, 1988.
- Beutner G., Sharma V.K., Giovannucci D.R., Yule D.I. and Sheu S-S. Identification of a ryanodine receptor in rat heart mitochondria. *J. Biol. Chem.* **276**, 21482-21488, 2001.
- Beutner G., Sharma V.K., Lin L., Ryu S-Y., Dirksen R.T. and Sheu S-S. Type 1 ryanodine receptor in cardiac mitochondria: Transducer of excitation-metabolism coupling. *Biochim. Biophys. Acta*, **1717**, 1-10, 2005.
- Bevan D.R. Neuromuscular relaxants. *Can. J. Anesth.*, **44**, 1135-1137, 1997.
- Bielawski J.P., Dunn K.A. and Yang Z. Rates of nucleotide substitution and mammalian nuclear gene evolution: Approximate and maximum-likelihood methods lead to different conclusions. *Genetics*, **156**, 1299-1308, 2000.
- Boerman R.H., Ophoff R.A., Links T.P., van Eijk R., Sandkuijl L.A., Elbaz A., Vale-Santos J.E., Wintzen A.R., van Deutekom J.C., Isles D.E., Fontaine B., Padberg G.W. and Frants R.R. Mutation in DHP receptor α_1 -subunit (CACLN1A3) gene in a Dutch family with hypokalaemic periodic paralysis. *J. Med. Genet.*, **32**, 44-47, 1995.
- Brandt A., Schleithoff L., Jurkat-Rott K., Klingler W., Baur C. and Lehmann-Horn F. Screening of the ryanodine receptor gene in 105 malignant hyperthermia families: Novel mutations and concordance with the *in vitro* contracture test. *Hum. Mol. Genet.*, **8**, 2055-2062, 1999.
- Brenig B. and Brem G. Genomic organisation and analysis of the 5' end of the porcine ryanodine receptor gene (*ryr1*). *FEBS Lett.*, **298**, 277-279, 1992.
- Breslauer K.J., Frank R., Blöcker H. and Marky L.A. Predicting DNA duplex stability from the base sequence. *Proc. Natl. Acad. Sci. USA.*, **83**, 3746-3750, 1986.
- Brilliantes A-M.B., Ondriaš K., Scott A., Kobrinsky E., Ondriašová E., Moschella M.C., Jayaraman T., Landers M., Ehrlich B.E. and Marks A.R. Stabilisation of calcium release channel (ryanodine receptor) function by FK506-binding protein. *Cell*, **77**, 513-523, 1994.
- Britt B.A. and Kalow W. Malignant hyperthermia: A statistical review. *Canad. Anaesth. Soc. J.*, **17**, 293-315, 1970.
- Britt B.A., Endrenyi L., Kalow W. and Peters P.L. The adenosine triphosphate (ATP) depletion test: Comparison with the caffeine contracture test as a method of diagnosing malignant hyperthermia susceptibility. *Can. Anaesth. Soc. J.*, **23**, 624-635, 1976.

- Britt B.A. Dantrolene. *Can. Anaesth. Soc. J.*, **31**, 61-75, 1984.
- Bronsveld I., Mekus F., Bijman J., Ballmann M., de Jonge H.R., Laabs U., Halley D.J., Ellemunter H., Mastella G., Thomas S., Veeze H.J. and Tümmler B. Chloride conductance and genetic background modulate the cystic fibrosis phenotype of $\Delta F508$ homozygous twins and siblings. *J. Clin. Invest.*, **108**, 1705-1715, 2001.
- Brook J.D., Harley H.G., Walsh K.V., Rundel S.A., Siciliano M.J., Harper P.S. and Shaw D.J. Identification of new DNA markers close to the myotonic dystrophy locus. *J. Med. Genet.*, **28**, 84-88, 1991.
- Brooksbank R.L., Badenhorst M.E., Isaacs H. and Savage N. Treatment of normal skeletal muscle with FK506 or rapamycin results in halothane-induced muscle contracture. *Anesthesiology*, **89**, 693-698, 1998.
- Brown R.L., Pollock N.A., Couchman K.G., Hodges M., Hutchinson D.O., Waaka R., Lynch P., McCarthy T.V. and Stowell K.M. A novel ryanodine receptor mutation and genotype-phenotype correlation in a large malignant hyperthermia New Zealand Maori pedigree. *Hum. Mol. Genet.*, **9**, 1515-1524, 2000.
- Brownell A.K.W., Paasuke R.T., Elash A., Fowlow S.B., Seagram C.G.F., Diewold R.J. and Friesen C. Malignant hyperthermia in Duchenne muscular dystrophy. *Anesthesiology*, **58**, 180-182, 1983.
- Brownell A.K.W. Malignant hyperthermia: Relationship to other diseases. *Br. J. Anaesth.*, **60**, 303-308, 1988.
- Brownie J., Shawcross S., Theaker J., Whitcombe D., Ferrie R., Newton C. and Little S. The elimination of primer-dimer accumulation in PCR. *Nucleic Acids Res.*, **25**, 3235-3241, 1997.
- Brunder D.G., Györke S., Dettbarn C. and Palade P. Involvement of sarcoplasmic reticulum Ca^{2+} release channels' in excitation-contraction coupling in vertebrate skeletal muscle. *J. Physiol.*, **445**, 759-778, 1992.
- Buratti R., Prestipino G., Menegazzi P., Treves S. and Zorzato F. Calcium dependent activation of skeletal muscle Ca^{2+} release channel (ryanodine receptor) by calmodulin. *Biochem. Biophys. Res. Commun.*, **213**, 1082-1090, 1995.
- Burns T.L., Moll P.P. and Schork M.A. Comparisons of different sampling designs for the determination of genetic transmission mechanisms in quantitative traits. *Am. J. Hum. Genet.*, **36**, 1060-1074, 1984.
- Callaway C., Seryshev A., Wang J-P., Slavik K.J., Needleman D.H., Cantu III C., Wu Y., Jayaraman T., Marks A.R. and Hamilton S.L. Localisation of the high and low affinity [3H]ryanodine binding sites on the skeletal muscle Ca^{2+} release channel. *J. Biol. Chem.*, **269**, 15879-15884, 1994.
- Capon F., Allen M.H., Ameen M., Burden D.A., Tillman D., Barker J.N. and Trembath R.C. A synonymous SNP of the corneodesmosin gene leads to increased mRNA stability and demonstrates association with psoriasis across diverse ethnic groups. *Hum. Mol. Genet.*, **13**, 2361-2368, 2004.
- Cargill M., Altshuler D., Ireland J., Sklar P., Ardlie K., Patil N., Lane C.R., Lim E.R., Kalyanaraman N., Nemesh J., Ziaugra L., Friedland L., Rolfe A., Warrington J., Lipshutz R., Daley G.Q. and Lander E.S. Characterisation of single-nucleotide polymorphisms in coding regions of human genes. *Nat. Genet.*, **22**, 231-238, 1999.
- Casana A., Fortunato G., De Sarno C., Brancadoro V. and Salvatore F. Identification of new polymorphisms in the CACNA1S gene. *Clin. Chem. Lab Med.*, **41**, 20-22, 2003.
- Catterall W.A. Structure and function of voltage-sensitive ion channels. *Science*, **242**, 50-61, 1988.
- Chamley D., Pollock N.A., Stowell K.M. and Brown R.L. Malignant hyperthermia in infancy and identification of novel RYR1 mutation. *Brit. J. Anaesth.*, **84**, 500-504, 2000.
- Chance B. and Williams G.R. Respiratory enzymes in oxidative phosphorylation.I. Kinetics of oxygen utilisation. *J. Biol. Chem.*, **22**, 383-393, 1955.
- Cheah K.S. and Cheah A.M. Mitochondrial calcium transport and calcium-activated phospholipase in porcine malignant hyperthermia. *Biochim. Biophys. Acta*, **634**, 70-84, 1981.
- Cheah K.S., Dauncey M.J., Cheah A.M. and Ingram D.L. Influence of environmental temperature and energy intake on porcine skeletal muscle mitochondria. *Comp. Biochem. Physiol.*, **82B**, 287-292, 1985a.
- Cheah K.S. and Cheah A.M. Malignant hyperthermia: Molecular defects in membrane permeability. *Experientia*, **41**, 656-661, 1985b.
- Cheah K.S., Chah A.M., Fletcher J.E. and Rosenberg H. Skeletal muscle mitochondrial respiration of malignant hyperthermia-susceptible patients. Ca^{2+} -induced uncoupling and free fatty acids. *Int. J. Biochem.*, **21**, 913-920, 1989.
- Chen S.R.W., Zhang L. and MacLennan D.H. Characterisation of a Ca^{2+} binding and regulatory site in the Ca^{2+} release channel (ryanodine receptor) of rabbit skeletal muscle sarcoplasmic reticulum. *J. Biol. Chem.*, **267**, 23318-23326, 1992.
- Chen S.R.W. and MacLennan D.H. Identification of calmodulin-, Ca^{2+} -, and ruthenium red-binding domains in the Ca^{2+} release channel (ryanodine receptor) of rabbit skeletal muscle sarcoplasmic reticulum. *J. Biol. Chem.*, **269**, 22698-22704, 1998.
- Chiang W., Allison C.P., Linz J.E. and Strasburg G.M. Identification of two α RYR alleles and characterisation of α RYR transcript variants in turkey skeletal muscle. *Gene*, **330**, 177-184, 2004.
- Chitayat D., Hodgkinson K.A., Ginsburg O., Dimmick J. and Waters G.V. King syndrome: A genetically heterogeneous phenotype due to congenital myopathies. *Am. J. Med. Genet.*, **43**, 954-956, 1992.

- Cho J.H., Nicolae D.L., Gold L.H., Fields C.T., LaBuda M.C., Rohal P.M., Pickles M.R., Qin L., Fu Y., Mann J.S., Kirschner B.S., Jabs E.W., Weber J., Hanauer S.B., Bayless T.M. and Brant S.R. Identification of novel susceptibility loci for inflammatory bowel disease on chromosomes 1p, 3q, and 4q: Evidence for epistasis between 1p and IBD1. *Proc. Natl. Acad. Sci. USA.*, **95**, 7502-7507, 1998.
- Christian A.S., Ellis F.R. and Halsall P.J. Is there a relationship between masseteric muscle spasm and malignant hyperpyrexia? *Br. J. Anaesth.*, **62**, 540-544, 1989.
- Chugun A., Taniguchi K., Murayama T., Uchide T., Hara Y., Temma K., Ogawa Y. and Akera T. Subcellular distribution of ryanodine receptors in the cardiac muscle of carp (*Cyprinus carpio*). *Am. J. Physiol. Regul. Integr. Comp. Physiol.*, **285**, R601-R609, 2003.
- Citino S.B., Bush M. and Phillips L.G. Dystocia and fatal hyperthermic episode in a giraffe. *J. Am. Vet. Med. Assoc.*, **185**, 1440-1442, 1984.
- Collins J.H., Tarcsafalvi A. and Ikemoto N. Identification of a region of calsequestrin that binds to the junctional face membrane of sarcoplasmic reticulum. *Biochem. Biophys. Res. Commun.*, **167**, 189-193, 1990.
- Collins J.S. and Schwartz C.E. Detecting polymorphisms and mutations in candidate genes. *Am. J. Hum. Genet.*, **71**, 1251-1252, 2002.
- Collins C.P. and Beirne R.O. Concepts in the prevention and management of malignant hyperthermia. *J. Oral Maxillofac. Surg.*, **61**, 1340-1345, 2003.
- Conti A., Gorza L. and Sorrentino V. Differential distribution of ryanodine type 3 (RyR3) gene product in mammalian skeletal muscles. *Biochem. J.*, **316**, 19-23, 1996.
- Cordell H.J. Epistasis: What it means, what it doesn't mean, and statistical methods to detect it in humans. *Hum. Mol. Genet.*, **11**, 2463-2468, 2002.
- Coronado R., Morrissette J., Sukhareva M. and Vaughan D.M. Structure and function of ryanodine receptors. *Am. J. Physiol.*, **266**, C1485-C1504, 1994.
- Crowder C.M., Shebester L.D. and Schedl T. Behavioural effects of volatile anesthetics in *Caenorhabditis elegans*. *Anesthesiology*, **85**, 901-912, 1996.
- Dalton D.L. Molecular screening for specific causative mutations in the South African malignant hyperthermia population. M.Sc. Thesis, North-West University, South Africa, 2004.
- Davies W., Harbitz I., Fries R., Stranzinger G. and Hauge J.G. Porcine malignant hyperthermia carrier detection and chromosomal assignment using a linked probe. *Anim. Genet.*, **19**, 203-212, 1988.
- Davis M., Brown R., Dickson A., Horton H., James D., Laing N., Marston R., Norgate M., Perlman D., Pollock N. and Stowell K. Malignant hyperthermia associated with exercise-induced rhabdomyolysis or congenital abnormalities and a novel RYR1 mutation in New Zealand and Australian pedigrees. *Br. J. Anaesth.*, **88**, 508-515, 2002.
- Davis M.R., Haan E., Jungbluth H., Sewry C., North K., Muntoni F., Kuntzer T., Lamont P., Bankier A., Tomlinson P., Sánchez A., Walsh P., Nagarajan L., Oley C., Colley A., Gedeon A., Quinlivan R., Dixon J., James D., Müller C.R. and Laing N.G. Principal mutation hotspot for central core disease and related myopathies in the C-terminal transmembrane region of the RYR1 gene. *Neuromuscul. Disord.*, **13**, 151-157, 2003.
- De Jong R.H., Heavner J.E. and Amory D.W. Malignant hyperpyrexia in the cat. *Anesthesiology*, **41**, 608-609, 1974.
- De La Vega F.M., Isaac H., Collins A., Scafe C.R., Halldórsson B.V., Su X., Lippert R.A., Wang Y., Laig-Webster M., Koehler R.T., Ziegler J.S., Wogan L.T., Stevens J.F., Leinen K.M., Olson S.J., Guegler K.J., You X., Xu L.H., Hemken H.G., Kalush F., Itakura M., Zheng Y., de Thé G., O'Brien S.J., Clark A.G., Istrail S., Hunkapiller M.W., Spier E.G. and Gilbert D.A. The linkage disequilibrium maps of three human chromosomes across four populations reflect their demographic history and a common underlying recombination pattern. *Genome Res.*, **15**, 454-462, 2005.
- Denborough M.A. and Lovell R.R.H. Anaesthetic deaths in a family. *Lancet*, **2**, 45, 1960.
- Denborough M.A., Foster J.F.A., Lovell R.R.H., Maplestone P.A. and Villiers J.D. Anaesthetic deaths in a family. *Brit. J. Anaesth.*, **34**, 395-396, 1962.
- Denborough M.A., Dennett X. and Anderson R. Central-core disease and malignant hyperpyrexia. *Br. Med. J.*, **1**, 272-273, 1973.
- Denborough M.A., Galloway G.J. and Hopkinson K.C. Malignant hyperthermia and sudden infant death. *Lancet*, **13**, 1068-1069, 1982.
- Denborough M. Malignant hyperthermia. *Lancet*, **352**, 1131-1136, 1998.
- Deufel T., Golla A., Iles D., Meindl A., Meitinger T., Schindelbauer D., DeVries A., Pongratz D., MacLennan D.H., Johnson K.J. and Lehmann-Horn F. Evidence of genetic heterogeneity of malignant hyperthermia susceptibility. *Am. J. Hum. Genet.*, **50**, 1151-1161, 1992.
- Deufel T., Sudbrak R., Feist Y., Rübsam B., Du Chesne I., Schäfer K.-L., Roewer N., Grimm T., Lehmann-Horn F., Hartung E. J. and Müller C. R. Discordance, in a malignant hyperthermia pedigree, between in vitro contracture-test phenotypes and haplotypes for the MHS1 region on chromosome 19q12-13.2, comprising the C1840T transition in the RYR1 gene. *Am. J. Hum. Genet.*, **56**, 1334-1342, 1995.

- Dickinson J.G. Heat-exercise hyperpyrexia. *J.R. Army Med. Corps.*, **135**, 27-29, 1989.
- Donati F. Muscle relaxants. A clinical update. *Can. J. Anesth.* **50**, R1-R4, 2003.
- Donnelly A.J. Malignant hyperthermia. *Assoc. Oper. Room Nurs.*, **59**, 393-405, 1994.
- Dorkins H.R. Suxamethonium – The development of a modern drug from 1906 to the present day. *Med. Hist.*, **26**, 145-168, 1982.
- Downey D. Porphyria: The road not traveled. *Med. Hypotheses*, **56**, 73-76, 2001.
- Du G.G., Sandhu B., Khanna V.K., Guo X.H. and MacLennan D.H. Topology of the Ca²⁺ release channel of skeletal muscle sarcoplasmic reticulum (RyR1). *Proc. Nat. Acad. Sci. USA.*, **26**, 16725-16730, 2002.
- Dubowitz V. and Pearse A.G.E. A comparative histochemical study of oxidative enzyme and phosphorylase activity in skeletal muscle. *Histochemie.*, **2**, 105-117, 1960.
- Ducart A., Adnet P., Renaud B., Riou B. and Krivosic-Horber R. Malignant hyperthermia during sevoflurane administration. *Anesth. Analg.*, **80**, 609-611, 1995.
- Ducreux S., Zorzato F., Ferreiro A., Jungbluth H., Muntoni F., Monnier N., Müller C.R. and Treves S. Functional properties of ryanodine receptors carrying 3 amino acid substitutions identified in patients affected by multi-minicore disease and central core disease, expressed in immortalised lymphocytes. *Biochem. J.*, **395**, 259-266, 2006.
- Dulhunty A.F. and Pouliquin P. What we don't know about the structure of ryanodine receptor calcium release channels. *Clin. Exp. Pharmacol. Physiol.*, **30**, 713-723, 2003.
- Dulhunty A.F., Karunasekara Y., Curtis S.M., Harvey P.J., Board P.G. and Casaraoito M.G. The recombinant dihydropyridine receptor II-III loop and partly structured 'C' region peptides modify cardiac ryanodine receptor activity. *Biochem. J.*, **385**, 803-813, 2005.
- Duret L. Evolution of synonymous codon usage in metazoans. *Curr. Opin. Genet. Dev.*, **12**, 640-649, 2002.
- Duthie G.G. and Arthur J.R. Free radicle and calcium homeostasis: Relevance to malignant hyperthermia? *Free Radical Bio. Med.*, **14**, 435-442, 1993.
- Eikelenboom G. and van den Bergh S.G. Mitochondrial metabolism in stress-susceptible pigs. *J. Anim. Sci.*, **37**, 692-696, 1973.
- Ellis F.R., Keaney N.P., Harriman D.G.F., Sumner D.W., Kyei-Mensah K., Tyrrell J.H., Hargreaves J.B., Parikh R.K. and Mulrooney P.L. Screening for malignant hyperpyrexia. *Br. Med. J.*, **3**, 559-561, 1972.
- Ellis F.R., Halsall P.J. and Christian A.S. Clinical presentation of suspected malignant hyperthermia during anaesthesia in 402 probands. *Anaesthesia*, **45**, 838-841, 1990.
- El-Orbany M.I., Joseph N.J., Salem M.R. and Klowden A.J. The neuromuscular effects and tracheal intubation conditions after small doses of succinylcholine. *Anesth. Analg.*, **98**, 1674-1675, 2004.
- Ervasti J.M., Claessens M.T., Mickelson J.R. and Louis C.F. Altered transverse tubule dihydropyridine receptor binding in malignant hyperthermia. *J. Biol. Chem.*, **264**, 2711-2717, 1989.
- European Malignant Hyperpyrexia Group. A protocol for the investigation of Malignant Hyperpyrexia (MH) susceptibility. *Br. J. Anaesth.*, **56**, 1267-1269, 1984.
- Eyre-Walker A. and Hurst L.D. The evolution of isochores. *Nature Rev. Genet.*, **2**, 549-555, 2001.
- Fagerlund T., Ørding H., Bendixen D. and Berg K. Search for three known mutations in the RYR1 gene in 48 Danish families with malignant hyperthermia. *Clin. Genet.*, **46**, 401-404, 1994.
- Fagerlund T., Ørding H., Bendixen D., Islander G., Ranklev-Twetman E. and Berg K. RYR mutation G1021A (Gly341Arg) is not frequent in Danish and Swedish families with malignant hyperthermia susceptibility. *Clin. Genet.*, **49**, 186-188, 1996.
- Fagerlund T.H., Ørding H., Bendixen D., Islander G., Ranklev-Twetman E. and Berg K. Discordance between malignant hyperthermia susceptibility and RYR1 mutation C1840T in two Scandinavian MH families exhibiting this mutation. *Clin. Genet.*, **52**, 416-421, 1997.
- Feil R. Environmental and nutritional effects on the epigenetic regulation of genes. *Mutat. Res.*, **600**, 46-57, 2006.
- Fessenden J.D., Feng W., Pessah I.N. and Allen P.D. Mutational analysis of putative calcium binding motifs within the skeletal muscle ryanodine receptor isoform RyR1. *J. Biol. Chem.*, **279**, 53028-53035, 2004.
- Feuerman T., Gade G.F. and Reynolds R. Stress-induced malignant hyperthermia in a head-injured patient. *J. Neurosurg.*, **68**, 297-299, 1988.
- Fiege M., Wappler F., Weissshorn R., Urich Gerbershagen M., Steinfath M. and Schulte am Esch J. Results of contracture tests with halothane, caffeine, and ryanodine depend on different malignant hyperthermia-associated ryanodine receptor gene mutations. *Anesthesiology*, **97**, 345-350, 2002.
- Filipski J. Correlation between molecular clock ticking, codon usage fidelity of DNA repair, chromosome banding and chromatin compactness in germline cells. *FEBS Lett.*, **217**, 184-186, 1987.
- Fill M., Coronado R., Mickelson J.R., Vilven J., Ma J., Jacobson B.A. and Louis C.F. Abnormal ryanodine receptor channels in malignant hyperthermia. *Biophys. J.*, **57**, 471-475, 1990.
- Fill M. and Copello J.A. Ryanodine receptor calcium release channels. *Physiol. Rev.*, **82**, 893-922, 2002.
- Flatt T. The evolutionary genetics of canalization. *Q. Rev. Biol.*, **80**, 287-316, 2005.

- Fleischer S., Ogunbunmi E.M., Dixon M.C. and Fleer E.A.M. Localisation of Ca²⁺ release channels with ryanodine in junctional terminal cisternae of sarcoplasmic reticulum of fast skeletal muscle. *Proc. Natl. Acad. Sci. USA.*, **82**, 7256-7259, 1985.
- Fletcher J.E., Tripolitis L., Erwin K., Hanson S. and Rosenberg H. Fatty acids modulate calcium-induced calcium release from skeletal muscle heavy sarcoplasmic reticulum fractions: Implications for malignant hyperthermia. *Biochem. Cell Biol.*, **68**, 1195-1201, 1990.
- Fletcher J.E., Conti P.A. and Rosenberg H. Comparison of North American and European Malignant Hyperthermia Group halothane contracture testing protocols in swine. *Acta Anaesthesiol. Scand.*, **35**, 483-487, 1991.
- Fletcher J.E., Calvo P.A. and Rosenberg H. Phenotypes associated with malignant hyperthermia susceptibility in swine genotyped as homozygous or heterozygous for the ryanodine receptor mutation. *Br. J. Anaesth.*, **71**, 410-417, 1993.
- Fletcher J.E., Tripolitis L., Hubert M., Vita G.M., Levitt R.C. and Rosenberg H. Genotype and phenotype relationships for mutations in the ryanodine receptor in patients referred for diagnosis of malignant hyperthermia. *Br. J. Anaesth.*, **75**, 307-310, 1995.
- Fletcher J.E., Wieland S.J., Karan S.M., Beech J. and Rosenberg H. Sodium channel in human malignant hyperthermia. *Anesthesiology*, **86**, 1023-1032, 1997.
- Fletcher J.E., Rosenberg H. and Aggarwal M. Comparison of European and North American malignant hyperthermia diagnostic protocol outcomes for use in genetic studies. *Anesthesiology*, **90**, 654-661, 1999.
- Fontaine B., Khurana T.S., Hoffman E.P., Bruns G.A.P., Haines J.L., Trofater J.A., Hanson M.P., Rich J., McFarlane H., Yasek D.M., Romano D., Gusella J.F. and Brown R.H. Hyperkalemic periodic paralysis and the adult muscle sodium channel α -subunit gene. *Science*, **250**, 1000-1002, 1990.
- Forrest K.A.T. and Cole P.J. Metabolic crises in anaesthesia. *Curr. Anaesth. Crit. Care*, **14**, 24-31, 2003.
- Fortunato G., Carsana A., Tinto N., Brancadoro V., Canfora G. and Salvatore F. A case of discordance between genotype and phenotype in a malignant hyperthermia family. *Eur. J. Hum. Genet.*, **7**, 415-420, 1999.
- Fortunato G., Berruti R., Brancadoro V., Fattore M., Salvatore F. and Casana A. Identification of a novel mutation in the ryanodine receptor gene (RYR1) in a malignant hyperthermia Italian family. *Eur. J. Hum. Genet.*, **8**, 149-152, 2000.
- Frayn K.N., Coppack S.W., Fielding B.A. and Humphreys S.M. Coordinated regulation of hormone-sensitive lipase and lipoprotein lipase in human adipose tissue *in vivo*: Implications for the control of fat storage and fat mobilization. *Advan. Enzyme Regul.*, **35**, 163-178, 1995.
- Freyssenet D., Irrcher I., Connor M.K., Di Carlo M. and Hood D.A. Calcium-regulated changes in mitochondrial phenotype in skeletal muscle cells. *Am. J. Physiol. Cell Physiol.*, **286**, C1053-C1061, 2004.
- Fruen B.R., Mickelson J.R. and Louis C.F. Dantrolene inhibition of sarcoplasmic reticulum Ca²⁺ release by direct and specific action at skeletal muscle ryanodine receptors. *J. Biol. Chem.*, **272**, 26965-26971, 1997.
- Fujii J., Otsu K., Zorzato F., De Leon S., Khanna V.K., Weiler J.E., O'Brien P.J. and MacLennan D.H. Identification of a mutation in porcine ryanodine receptor associated with malignant hyperthermia. *Science*, **253**, 448-451, 1991.
- Fullerton S.M., Carvalho A.B. and Clark A.G. Local rates of recombination are positively correlated with GC content in the human genome. *Mol. Biol. Evol.*, **18**, 1139-1142, 2001.
- Furrer B., Candrian U., Wieland P. and Lüthy J. Improving PCR efficiency. *Nature*, **346**, 324, 1990.
- Fusi F., Iozzi D., Sgaragli G. and Frosini M. 3,5-di-*t*-butylcatechol (DTCAT) as an activator of rat skeletal muscle ryanodine receptor Ca²⁺ channel (RyRC). *Biochem. Pharmacol.*, **69**, 485-491, 2005.
- Futatsugi A., Kuwajima G. and Mikoshiba K. Tissue-specific and developmentally regulated alternative splicing in mouse skeletal muscle ryanodine receptor mRNA. *Biochem. J.*, **305**, 373-378, 1995.
- Gallant E.M. and Rempel W.E. Porcine malignant hyperthermia: False negatives in the halothane test. *Am. J. Vet. Res.*, **48**, 488-491, 1987.
- Gallen J.S. Propofol does not trigger malignant hyperthermia. *Anesth. Analg.*, **72**, 406-416, 1991.
- Galli L., Orrico A., Cozzolino S., Pietrini V., Tegazzin V. and Sorrentino V. Mutations in the RYR1 gene in Italian patients at risk for malignant hyperthermia: Evidence for a cluster of novel mutations in the C-terminal region. *Cell Calcium*, **32**, 143-151, 2002.
- Galli L., Orrico A., Lorenzini S., Censini S., Falciani M., Covacci A., Tegazzin V. and Sorrentino V. Frequency and localisation of mutations in the 106 exons of the RYR1 gene in 50 individuals with malignant hyperthermia. *Hum. Mutat.*, **27**, 830-838, 2006.
- Galloway G.J. and Denborough M.A. Suxamethonium chloride and malignant hyperpyrexia. *Br. J. Anaesth.*, **58**, 447-450, 1986.
- Garrido S., Fraga M., Martin M.J. and Belda J. Malignant hyperthermia during desflurane-succinylcholine anesthesia for orthopedic surgery. *Anesthesiology*, **90**, 1208-1209, 1999.
- George A.L., Ledbetter D.H., Kallen R.G. and Barchi R.L. Assignment of a human skeletal muscle sodium channel α -subunit gene (SCN4A) to 17q23.1-25.3. *Genomics*, **9**, 555-556, 1991.

- George C.H., Jundi H., Thomas N.L., Scoote M., Walters N., Williams A.J. and Lai F.A. Ryanodine receptor regulation by intramolecular interaction between cytoplasmic and transmembrane domains. *Mol. Biol. Cell*, **6**, 2627-2638, 2004.
- Giannini G., Clementi E., Ceci R., Marziali G. and Sorrentino V. Expression of a ryanodine receptor-Ca²⁺ channel that is regulated by TGF- β . *Science*, **257**, 91-94, 1992.
- Giannini G.H., Conti A., Mammarella S., Scrobogna M. and Sorrentino V. The ryanodine receptor/calcium channel genes are widely and differentially expressed in murine brain and peripheral tissues. *J. Cell Biol.*, **128**, 893-904, 1995.
- Gillard E.F., Otsu K., Fujii J., Khanna V.K., De Leon S., Derdemezi J., Britt B.A., Duff C.L., Worton R.G. and MacLennan D.H. A substitution of cysteine for arginine 614 in the ryanodine receptor is potentially causative of human malignant hyperthermia. *Genomics*, **11**, 751-755, 1991.
- Gillard E.F., Otsu K., Fujii J., Duff C., De Leon S., Khanna V.K., Britt B.A., Worton R.G. and MacLennan D.H. Polymorphisms and deduced amino acid substitutions in the coding sequence of the ryanodine receptor (RYR1) gene in individuals with malignant hyperthermia. *Genomics*, **13**, 1247-1254, 1992.
- Gilly H., Musat I., Fricker R., Bittner R.E., Steinbereithner K. and Kress H-G. Classification of malignant hyperthermia-equivocal patients by 4-chloro-*m*-cresol. *Anesth. Analg.*, **85**, 149-154, 1997.
- Ginsburg R. and Purcell-Jones G. Malignant hyperthermia in the Wolf-Hirschhorn syndrome. *Anaesthesia*, **43**, 386-388, 1988.
- Girard T., Urwyler A., Censier K., Mueller C.R., Zorzato F. and Treves S. Genotype-phenotype comparison of the Swiss malignant hyperthermia population. *Hum. Mutat.*, **18**, 357-358, 2001.
- Girard T., Treves S., Voronkov E., Siegemund M. and Urwyler A. Molecular genetic testing for malignant hyperthermia susceptibility. *Anesthesiology*, **100**, 1076-1080, 2004.
- Golinski M. Malignant hyperthermia: A review. *Plast. Surg. Nurs.*, **15**, 30-34, 1995.
- Gonzalez L.E., Meléndez-Vásquez C.V., Gregson N.A. and File S.E. A rat model of spontaneous myopathy and malignant hyperthermia. *Am. J. Pathol.*, **152**, 1099-1103, 1998.
- Grinberg R., Edelstein G. and Gordon A. Postoperative malignant hyperthermia episodes in patients who received "safe" anaesthetics. *Can. Anaesth. Soc. J.*, **30**, 273-276, 1983.
- Groenewald J.Z., Liebenberg J., Groenewald I.M. and Warnich L. Linkage disequilibrium analysis in a recently founded population: Evaluation of the variegated porphyria founder in South African Afrikaners. *Am. J. Hum. Genet.*, **62**, 1254-1258, 1998.
- Groh S., Marty I., Ottolia M., Prestipino G., Chapel A., Villaz M. and Ronjat M. Functional interaction of the cytoplasmic domain of triadin with the skeletal ryanodine receptor. *J. Biol. Chem.*, **274**, 12278-12283, 1999.
- Gronert G.A. Malignant hyperthermia. *Anesthesiology*, **53**, 395-423, 1980.
- Guis S., Figarella-Branger D., Monnier N., Bendahan D., Kozak-Ribbens G., Mattei J-P., Lunardi J., Cozzone P.J. and Pellissier J-P. Multiminicore disease in a family susceptible to malignant hyperthermia. *Arch. Neurol.*, **61**, 106-113, 2004.
- Guo W. and Campbell K.P. Association of triadin with the ryanodine receptor and calsequestrin in the lumen of the sarcoplasmic reticulum. *J. Biol. Chem.*, **270**, 9027-9030, 1995.
- Guo Y. and Jamison D.C. The distribution of SNPs in human gene regulatory regions. *BMC Genomics*, **6**, 140-151, 2005.
- Gurrera R.J. Is neuroleptic malignant syndrome a neurogenic form of malignant hyperthermia? *Clin. Neuropharmacol.*, **25**, 183-193, 2002.
- Hackl W., Mauritz W., Schemper M., Winkler M., Sporn P. and Steinbereithner K. Prediction of malignant hyperthermia susceptibility: Statistical evaluation of clinical signs. *Br. J. Anaesth.*, **64**, 425-429, 1990.
- Hakamata Y., Nishimura S., Nakai J., Nakashima Y., Kita T. and Imoto K. Involvement of the brain type of ryanodine receptor in T-cell proliferation. *FEBS Lett.*, **352**, 206-210, 1994.
- Hall L.W., Woolf N., Bradley J.W. and Jolly D.W. Unusual reaction to suxamethonium chloride. *Br. Med. J.*, **2**, 1305, 1966.
- Hall T.A. BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucl. Acids Symp. Ser.*, **41**, 95-98, 1999.
- Hall-Curran J.L., Stewart A.D., Ball S.P., Halsall J.P., Hopkins P.M. and Ellis F.R. No C1840 to T in RYR1 in malignant hyperthermia. *Hum. Mutat.*, **2**, 330, 1993.
- Halsall P.J., Cain P.A. and Ellis F.R. Retrospective analysis of anaesthetics received by patients before susceptibility to malignant hyperpyrexia was recognised. *Br. J. Anaesth.*, **51**, 949-954, 1979.
- Halsall P.J. and Ellis F.R. Malignant hyperthermia in Baillière's Clinical anaesthesiology: International practice and research. W.B. Saunders, London, UK. **7**, 343-356, 1993.
- Halsall J. and Robinson R. Genetic testing for malignant hyperthermia. *Curr. Anaesth. Crit. Care*, **15**, 11-21, 2004.
- Hans M., Luviset S., Williams M.E., Spagnolo M., Urrutia A., Tottene A., Brust P.F., Johnson E.C., Harpold M.M., Stauderman K.A. and Pietrobon D. Functional consequences of mutations in the human α_{1A} calcium channel subunit linked to familial hemiplegic migraine. *J. Neurosci.*, **19**, 1610-1619, 1999.

- Harbitz I., Chowdhary B., Thomsen P.D., Davies W., Kaufmann U., Kran S., Gustavsson I., Christensen K. and Hauge J.G. Assignment of the porcine calcium release channel gene, a candidate for the malignant hyperthermia locus, to the 6p11-q21 segment of chromosome 6. *Genomics*, **8**, 243-248, 1990.
- Hare M.P. and Palumbi S.R. High intron sequence conservation across three mammalian orders suggests functional constraints. *Mol. Biol. Evol.*, **20**, 969-978, 2003.
- Harrison G.G., Saunders S.J., Biebuyck J.F., Hickman R., Dent D.M., Weaver V. and Terblanche J. Anaesthetic-induced malignant hyperpyrexia and a method for its prediction. *Br. J. Anaesth.*, **41**, 844-854, 1969.
- Harrison G.G. Dantrolene sodium in the treatment of malignant hyperthermia. *S. Afr. Med. J.*, **5**, 909, 1981.
- Harrison G.G. and Isaacs H. Malignant hyperthermia. An historical vignette. *Anaesthesia*, **47**, 54-56, 1992.
- Hartung E., Koob M., Anetseder M., Schoemig P., Krauspe R., Hogrefe G. and Engelhardt W. Malignant hyperthermia (MH) diagnostics: A comparison between the halothane-caffeine- and the ryanodine-contracture-test results in MH susceptibility, normal and control muscle. *Acta Anaesthesiol. Scand.*, **40**, 437-444, 1996.
- Havenga Y. Screening for specific mutations in malignant hyperthermia susceptible families. M.Sc. Thesis, University of Pretoria, South Africa, 2000.
- Hawkes M.J., Nelson T. and Hamilton S.L. [³H]Ryanodine as a probe of changes in the functional state of the Ca²⁺-release channel in malignant hyperthermia. *J. Biol. Chem.*, **267**, 6702-6709, 1992.
- Hayashi K., Miller R.G. and Brownell K.W. Central core disease: Ultrastructure of the sarcoplasmic reticulum and t-tubules. *Muscle Nerve*, **12**, 95-102, 1989.
- Hayek S.M., Zhu X., Bhat M.B., Zhao J., Takeshima H., Valdivia H.H. and Ma J. Characterisation of a calcium-regulation domain of the skeletal-muscle ryanodine receptor. *Biochem. J.*, **351**, 57-65, 2000.
- Healy J.M.S., Quane K.A., Keating K.K., Lehane M., Heffron J.J.A. and McCarthy T.V. Diagnosis of malignant hyperthermia: A comparison of the *in vitro* contracture test with the molecular genetic diagnosis in a large pedigree. *J. Med. Genet.*, **33**, 18-24, 1996.
- Hellmann I., Zöllner S., Enard W., Ebersberger I., Nickel B. and Pääbo S. Selection on human genes as revealed by comparisons to chimpanzee cDNA. *Genome Res.*, **13**, 831-837, 2003.
- Hermesh H., Aizenberg D., Lapidot M. and Munitz H. Risk of malignant hyperthermia among patients with neuroleptic malignant syndrome and their families. *Am. J. Psychiatry*, **145**, 1431-1434, 1988.
- Herrmann M.G., Durtschi J.D., Bromley L.K., Witter C.T. and Voelkerding K.V. Amplicon DNA melting analysis for mutation scanning and genotyping: Cross-platform comparison of instruments and dyes. *Clin. Chem.*, **52**, 494-503, 2006.
- Heytens L., Martin J.J., van de Kelft E. and Bossaert L.L. *In vitro* contracture tests in patients with various neuromuscular diseases. *Br. J. Anaesth.*, **68**, 72-75, 1992.
- Hidalgo C., Bull R., Behrens M.I. and Donoso P. Redox regulation of RyR-mediated Ca²⁺ release in muscle and neurons. *Biol. Res.*, **37**, 716-760, 2004.
- Hogan K., Couch F., Powers P.A. and Gregg R.G. A cysteine-for-arginine substitution (R614C) in the human skeletal muscle calcium release channel cosegregates with malignant hyperthermia. *Anesth. Analg.*, **75**, 441-448, 1992.
- Hogan K. The anesthetic myopathies and malignant hyperthermias. *Curr. Opin. Neurol.*, **11**, 469-476, 1998.
- Holliday R. A different kind of inheritance. *Sci. Am.*, **1**, 60-73, 1989.
- Holliday R. Epigenetics comes of age in the twenty first century. *J. Genet.*, **81**, 1-4, 2002.
- Hopkins P.M., Ellis F.R. and Halsall P.J. Ryanodine contracture: A potentially specific *in vitro* diagnostic test for malignant hyperthermia. *Br. J. Anaesth.*, **66**, 611-613, 1991.
- Hopkins P.M., Ellis F.R., Halsall P.J. and Stewart A.D. An analysis of the predictive probability of the *in vitro* contracture test for determining susceptibility to malignant hyperthermia. *Anesth. Analg.*, **84**, 648-656, 1997.
- Hopkins P.M. Malignant hyperthermia: Advances in clinical management and diagnosis. *Br. J. Anaesth.*, **85**, 118-128, 2000.
- Hosoi E., Nishizaki C., Gallagher K.L., Wyre H.W., Matsuo Y. and Sei Y. Expression of the ryanodine receptor isoforms in immune cells. *J. Immunol.*, **167**, 4887-4894, 2001.
- Huang C-H., Reid M., Daniels G. and Blumenfeld O.O. Alteration of splice site selection by an exon mutation in the human glycophorin A gene. *J. Biol. Chem.*, **268**, 25902-25908, 1993.
- Hubbard T., Andrews D., Caccamo M., Cameron G., Chen Y., Clamp M., Clarke L., Coates G., Cox T., Cunningham F., Curwen V., Cutts T., Down T., Durbin R., Fernandez-Suarez X.M., Gilbert J., Hammond M., Herrero J., Hotz H., Howe K., Iyer V., Jekosch K., Kahari A., Kasprzyk A., Keefe D., Keenan S., Kokocinski F., London D., Longden I., McVicker G., Melsopp C., Meidl P., Potter S., Proctor G., Rae M., Rios D., Schuster M., Searle S., Severin J., Slater G., Smedley D., Smith J., Spooner W., Stabenau A., Stalker J., Storey R., Trevanion S., Ureta-Vidal A., Vogel J., White S., Woodwark C. and Birney E. Ensemble 2005. *Nucleic Acids Res.*, **33**, D447-D453, 2005.
- Iaizzo P.A., Seewald M., Oakes S.G. and Lehmann-Horn F. The use of Fura-2 to estimate myoplasmic (Ca²⁺) in human skeletal muscle. *Cell Calcium*, **10**, 151-158, 1989.

- Iaizzo P.A. and Lehmann-Horn F. Anaesthetic complications in muscle disorders. *Anesthesiology*, **82**, 1093-1096, 1995.
- Ibarra C.A., Wu S., Murayama K., Minami N., Ichihara Y., Kikuchi H., Noguchi S., Hayashi Y.K., Ochiai R. and Nishino I. Malignant hyperthermia in Japan. Mutation screening of the entire ryanodine receptor type 1 gene coding region by direct sequencing. *Anesthesiology*, **104**, 1146-1154, 2006.
- Ikemoto N., Bhatnagar G.M., Nagy B. and Gergely J. Interaction of divalent cations with the 55,000-dalton protein component of the sarcoplasmic reticulum. Studies of fluorescence and circular dichroism. *J. Biol. Chem.*, **247**, 7835-7837, 1972.
- Iles D.E., Segers B., Heytens L., Sengers R.C.A and Wieringa B. High-resolution physical mapping of four microsatellite repeat markers near the RYR1 locus on chromosome 19q13.1 and apparent exclusion of the MHS locus from this region in two malignant hyperthermia susceptible families. *Genomics*, **14**, 749-754, 1992.
- Iles D.E., Segers B., Sengers R.C.A., Monsieurs K., Heytens L., Halsall P.J., Hopkins P.M., Ellis F.R., Hall-Curran J.L., Stewart A.D. and Wieringa B. Genetic mapping of the β_1 - and γ -subunits of the human skeletal muscle L-type voltage-dependent calcium channel on chromosome 17q and exclusion as candidate genes for malignant hyperthermia. *Hum. Mol. Genet.*, **2**, 863-868, 1993.
- Iles D.E., Lehmann-Horn F., Scherer S.W., Tsui L-C., Olde Weghuis D., Suijkerbuijk R.F., Heytens L., Mikala G., Schwartz A., Ellis F.R., Stewart A.D., Deufel T. and Wieringa B. Localisation of the gene encoding the α_2/δ -subunits of the L-type voltage-dependant calcium channel to chromosome 7q and analysis of the segregation of flanking markers in malignant hyperthermia susceptible families. *Hum. Mol. Genet.*, **3**, 969-975, 1994.
- Imagawa T., Smith J.S., Coronado R. and Campbell K.P. Purified ryanodine receptor from skeletal muscle sarcoplasmic reticulum is the Ca^{2+} -permeable pore of the calcium release channel. *J. Biol. Chem.*, **262**, 16636-16643, 1987.
- Imbert N., Cognard C., Dupont G., Guillou C. and Raymond G. Abnormal calcium homeostasis in Duchenne muscular dystrophy myotubes contracting *in vitro*. *Cell Calcium*, **18**, 177-186, 1995.
- Isaacs H., Heffron J.J.A. and Badenhorst M. Predictive test for malignant hyperpyrexia. *Br. J. Anaesth.*, **47**, 1075-1080, 1975.
- Isaacs H. and Gericke G. Concurrence of malignant hyperthermia and congenital abnormalities. *Muscle Nerve*, **13**, 915-917, 1990.
- Isaacs H. and Badenhorst M. False-negative results with muscle caffeine halothane contracture testing for malignant hyperthermia. *Anesthesiology*, **79**, 5-9, 1993.
- Islander G., Henriksson K-G. and Ranklev-Twetman E. Malignant hyperthermia susceptibility without central core disease (CCD) in a family where CCD is diagnosed. *Neuromusc. Disord.*, **5**, 125-127, 1995.
- Islander G. and Twetman E.R. Comparison between the European and North American protocols for diagnosis of malignant hyperthermia susceptibility in humans. *Anesth. Analg.*, **88**, 1155-1160, 1999.
- Jaenisch R. and Bird A. Epigenetic regulation of gene expression: How the genome integrates intrinsic and environmental signals. *Nat. Genet.*, **33**, 245-254, 2003.
- Jay S.D., Ellis S.B., McCue A.F., Williams M.E., Vedvick T.S., Harpold M.M. and Campbell K.P. Primary structure of the γ -subunit of the DHP-sensitive calcium channel from skeletal muscle. *Science*, **248**, 490-491, 1990.
- Jayaraman T., Brillantes A-M., Timerman A.P., Fleischer S., Erdjument-Bromage H., Tempst P. and Marks A.R. FK506 binding protein associated with the calcium release channel (ryanodine receptor). *J. Biol. Chem.*, **267**, 9474-9477, 1992.
- Jiang D., Xiao B., Li X. and Chen S.R.W. Smooth muscle tissues express a major dominant negative splice variant of the type 3 Ca^{2+} release channel (ryanodine receptor). *J. Biol. Chem.*, **278**, 4763-4769, 2003.
- Jungbluth H., Zhou H., Bertini E., Treves S., Sewry C. and Muntoni F. Autosomal dominant mutations in the human ryanodine receptor (RYR1) gene associated with centronuclear myopathy. *Neuromuscul. Disord.*, **15**, 680, 2005.
- Jurkat-Rott K., Lehmann-Horn F., Elbaz A., Heine R., Gregg R.G., Hogan K., Powers P.A., Lapie P., Vale-Santos J.E., Weissenbach J. and Fontaine B. A calcium channel mutation causing hypokalemic periodic paralysis. *Hum. Mol. Genet.*, **3**, 1415-1419, 1994.
- Jurkat-Rott K., McCarthy T. and Lehmann-Horn F. Genetics and pathogenesis of malignant hyperthermia. *Muscle Nerve*, **23**, 4-17, 2000.
- Jurkat-Rott K. and Lehmann-Horn F. Muscle channelopathies and critical points in functional and genetic studies. *J. Clin. Invest.*, **115**, 2000-2009, 2005.
- Kalow W., Britt B.A., Terreau M.E. and Haist C. Metabolic error of muscle metabolism after recovery from malignant hyperthermia. *Lancet*, **7679**, 895-898, 1970.
- Katsuya H., Misumi M., Ohtani Y. and Miike T. Postanaesthetic acute renal failure due to carnitine palmityl transferase deficiency. *Anesthesiology*, **68**, 945-948, 1988.
- Kaus S.J. and Rockoff M.A. Malignant hyperthermia. *Pediatr. Anaesth.*, **41**, 221-236, 1994.

- Kausch K., Lehmann-Horn F., Janka M., Wieringa B., Grimm T. and Müller C.R. Evidence for linkage of the central core disease locus to the proximal long arm of human chromosome 19. *Genomics*, **10**, 765-769, 1991.
- Keating K.E., Quane K.A., Manning B.M., Lehane M., Hartung E., Censier K., Urwyler A., Klausnitzer M., Muller C.R., Heffron J.J.A. and McCarthy T.V. Detection of a novel RYR1 mutation in four malignant hyperthermia pedigrees. *Hum. Mol. Genet.*, **3**, 1855-1858, 1994.
- Keating K.E., Giblin L., Lynch P.J., Quane K.A., Lehane M., Heffron J.J.A. and McCarthy T.V. Detection of a novel mutation in the ryanodine receptor gene in an Irish malignant hyperthermia pedigree: Correlation of the IVCT response with the affected and unaffected haplotypes. *J. Med. Genet.*, **34**, 291-296, 1997.
- Keith J.M., Cochran D.A.E., Lala G.H., Adams P., Bryant D. and Mitchelson K.R. Unlocking hidden genomic sequence. *Nucleic Acids Res.*, **32**, e35, 2004.
- Kelfer H.M., Singer W.D. and Reynolds R.N. Experience and reason—Briefly recorded. *Pediatrics*, **71**, 118-119, 1983.
- Kim C.H., Oh Y. and Lee T.H. Codon optimisation for high-level expression of human erythropoietin (EPO) in mammalian cells. *Gene*, **199**, 293-301, 1997.
- King J.O., Denborough M.A. and Zapf P.W. Inheritance of malignant hyperpyrexia. *Lancet*, **12**, 365-370, 1972.
- Kirichok Y., Krapivinsky G. and Clapham D.E. The mitochondrial calcium uniporter is a highly selective ion channel. *Nature*, **427**, 360-364, 2004.
- Klein L.V. Case report: A hot horse. *Vet. Anesth.*, **2**, 41-42, 1975.
- Kleopa K.A., Rosenberg H. and Heiman-Patterson T. Malignant hyperthermia-like episode in Becker muscular dystrophy. *Anesthesiology*, **93**, 1535-1537, 2000.
- Klingler W., Baur C., Georgieff M., Lehmann-Horn F. and Melzer W. Detection of proton release from cultured human myotubes to identify malignant hyperthermia susceptibility. *Anesthesiology*, **97**, 1059-1066, 2002.
- Knudson M.C., Chaudhari N., Sharp A.H., Powell J.A., Beam K.G. and Campbell K.P. Specific absence of the α_1 -subunit of the dihydropyridine receptor in mice with muscular dysgenesis. *J. Biol. Chem.*, **264**, 1345-1348, 1989.
- Kobayashi S., Yamamoto T., Parness J. and Ikemoto N. Antibody probe study of Ca^{2+} channel regulation by interdomain interaction within the ryanodine receptor. *Biochem. J.*, **380**, 561-569, 2004.
- Kolb M.E., Horne M.L. and Martz R. Dantrolene in human malignant hyperthermia: A multicenter study. *Anaesthesiology*, **56**, 254-262, 1982.
- Korch C. and Drabkin H. Improved DNA sequencing accuracy and detection of heterozygous alleles using manganese citrate and different fluorescent dye terminators. *Genome Res.*, **9**, 588-595, 1999.
- Kossugue P.M., Muinz V.P., Pavanello R.C.M., Silva H.C., Gurgel Giannetti J., Paim J.F.O., Zatz M. and Vainzof M. Screening for mutations in the C-terminal region of the RYR1 gene identify high frequency of autosomal recessive form of central core disease (CCD) in Brazil. *Neuromuscul. Disord.*, **15**, 676-743, 2005.
- Kraus R.L., Sinnegger M.J., Koschak A., Glossmann H., Stenirri S., Carrera P. and Striessnig J. Three new familial hemiplegic migraine mutants affect P/Q-type Ca^{2+} channel kinetics. *J. Biol. Chem.*, **275**, 9239-9243, 2000.
- Krueger G. and Ellis C.N. Psoriasis. Recent advances in understanding its pathogenesis and treatment. *J. Am. Acad. Dermatol.*, **53**, S94-100, 2005.
- Kruglyak L. The use of a genetic map of biallelic markers in linkage studies. *Nat. Genet.*, **17**, 21-24, 1997.
- Kubota T., Horie M., Takano M., Yoshida H., Takenaka K., Watanabe E., Tsuchiya T., Otani H. and Sasayama S. Evidence for a single nucleotide polymorphism in the KCNQ1 potassium channel that underlies susceptibility to life-threatening arrhythmias. *J. Cardiovasc. Electrophysiol.*, **12**, 1223-1229, 2001.
- Kunst G., Graf B.M., Schreiner R., Martin E. and Fink R.H. Differential effects of sevoflurane, isoflurane, and halothane on Ca^{2+} release from the sarcoplasmic reticulum of skeletal muscle. *Anesthesiology*, **91**, 179-186, 1999.
- Kunst G., Stucke A.G., Graf B.M., Martin E. and Fink R.H.A. Desflurane induces only minor Ca^{2+} release from the sarcoplasmic reticulum of mammalian skeletal muscle. *Anesthesiology*, **93**, 832-839, 2000.
- Kuzmenkin A., Jurkat-Rott K., Lehmann-Horn F. and Mitrovic N. Impaired slow inactivation due to a polymorphism and substitutions of Ser-906 in the II-III loop of the human $\text{Na}_v1.4$ channel. *Eur. J. Physiol.*, **447**, 71-77, 2003.
- Laan M. and Pääbo S. Demographic history and linkage disequilibrium in human populations. *Nat. Genet.*, **17**, 435-438, 1997.
- Lai F.A., Misra M., Xu L., Smith A. and Meissner G. The ryanodine receptor- Ca^{2+} release channel complex of skeletal muscle sarcoplasmic reticulum. *J. Biol. Chem.*, **264**, 16776-16785, 1989.
- Laitinen P.J., Brown K.M., Piippo K., Swan H., Devaney J.M., Brahmhatt B., Donarum E.A., Marino M., Tiso N., Viitasalo M., Toivonen L., Stephan D.A., Kontula K., Laitinen P.J., Brown K.M., Piippo K., Swan

- H., Devaney J.M., Brahmabhatt B., Donarum E.A., Marino M., Tiso N., Viitasalo M., Toivonen L., Stephan D.A. and Kontula K. Mutations of the cardiac ryanodine receptor (RyR2) gene in familial polymorphic ventricular tachycardia. *Circulation*, **103**, 485-490, 2001.
- Larach M.G. for the North American Malignant Hyperthermia Group. Standardisation of the caffeine halothane muscle contracture test. *Anesth. Analg.*, **69**, 511-515, 1989.
- Larach M.G., Landis J.R., Shirk S.J., Diaz M. and the North American malignant hyperthermia registry. Prediction of malignant hyperthermia susceptibility in man: Improving sensitivity of the caffeine halothane contracture test. *Anesthesiology*, **77**, A1052, 1992.
- Larach M.G. Should we use muscle biopsy to diagnose malignant hyperthermia susceptibility? *Anesthesiology*, **79**, 1-4, 1993.
- Larach M.G., Localio A.R., Allen G.C., Denborough M.A., Ellis F.R., Gronert G.A., Kaplan R.F., Muldoon S.M., Nelson T.E., Ørding H., Rosenberg H., Waud B.E. and Wedel D.J. A clinical grading scale to predict malignant hyperthermia susceptibility. *Anesthesiology*, **80**, 771-779, 1994.
- Lawrie R.A. Post mortem glycolysis in normal and exudative longissimus dorsi muscles of the pig in relation to so-called white muscle disease. *J. Comp. Path.*, **70**, 243-295, 1960.
- Ledbetter M.W., Preiner J.K., Louis C.F. and Mickelson J.R. Tissue distribution of ryanodine receptor isoforms and alleles determined by reverse transcription polymerase chain reaction. *J. Biol. Chem.*, **269**, 31544-31551, 1994.
- Lee B.S., Sessanna S., Laychock S.S. and Rubin R.P. Expression and cellular localization of a modified type 1 ryanodine receptor and L-type channel proteins in non-muscle cells. *J. Membrane Biol.*, **189**, 181-190, 2002.
- Lees D.E., Gadde P.L. and Macnamara T.E. Malignant hyperthermia in association with Burkitt's lymphoma: Report of a third case. *Anesth. Analg.*, **59**, 514-515, 1980.
- Leong P. and MacLennan D.H. The cytoplasmic loops between domains II and III and domains III and IV in the skeletal muscle dihydropyridine receptor bind to a contiguous site in the skeletal muscle ryanodine receptor. *J. Biol. Chem.*, **273**, 29958-29964, 1998.
- Levitt R.C., McKusick V.A., Fletcher J.E. and Rosenberg H. Gene candidate. *Nature*, **345**, 297-298, 1990.
- Levitt R.C., Nouri N., Jedlicka A.E., McKusick V.A., Marks A.R., Shutack J.G., Fletcher J.E., Rosenberg H. and Meyers D.A. Evidence for genetic heterogeneity in malignant hyperthermia susceptibility. *Genomics*, **11**, 543-547, 1991.
- Levitt R.C., Olickers A., Meyers S., Fletcher J.E., Rosenberg H., Isaacs H. and Meyers D.A. Evidence for the localisation of a malignant hyperthermia susceptibility locus (MHS2) to human chromosome 17q. *Genomics*, **14**, 562-566, 1992.
- Liu G. and Pessah I.N. Molecular interaction between ryanodine receptor and glycoprotein triadin involved redox cycling of functionally important hyperreactive sulfhydryls. *J. Biol. Chem.*, **269**, 33028-33034, 1994.
- Liu H.X., Cartegni L., Zhang M.Q. and Krainer A.R. A mechanism for exon skipping caused by nonsense or missense mutations in BRCA1 and other genes. *Nat. Genet.*, **27**, 55-58, 2001.
- Lo H.S., Wang Z., Hu Y., Yang H.H., Gere S., Buetow K.H. and Lee M.P. Allelic variation in gene expression is common in the human genome. *Genome Res.*, **13**, 1855-1862, 2003.
- Loghmanee F. and Tobak M. Fatal malignant hyperthermia associated with recreational cocaine and ethanol abuse. *Am. J. Forensic Med. Pathol.*, **7**, 246-248, 1986.
- Loke J. and MacLennan D.H. Malignant hyperthermia and central core disease: Disorders of Ca²⁺ release channels. *Am. J. Med.*, **104**, 470-486, 1998.
- Loke J.C.P., Kraev N., Sharma P., Du G., Patel L., Kreav A. and MacLennan D.H. Detection of a novel ryanodine receptor subtype 1 mutation (R328W) in a malignant hyperthermia family by sequencing of a leukocyte transcript. *Anesthesiology*, **99**, 297-302, 2003.
- Lopez J.R., Perez C., Linares N., Allen P. and Terzic A. Hypersensitive response of malignant hyperthermia-susceptible skeletal muscle to inositol 1,4,5-triphosphate induced release of calcium. *Naunyn Schmiedebergs Arch. Pharmacol.*, **352**, 442-446, 1995.
- Louis C.F., Zualkernan K., Roghair T. and Mickelson J.R. The effect of volatile anesthetics on calcium regulation by malignant hyperthermia. Susceptible sarcoplasmic reticulum. *Anesthesiology*, **77**, 114-125, 1992.
- Lu X., Xu L. and Meissner G. Activation of the skeletal muscle calcium release channel by a cytoplasmic loop of the dihydropyridine receptor. *J. Biol. Chem.*, **269**, 6511-6516, 1994.
- Luedtke N.W., Liu Q. and Tor Y. On the electronic structure of ethidium. *Chem. Eur. J.*, **7**, 495-508, 2005.
- Lusis A.J., Heinzmann C., Sparkes R.S., Scott J., Knott T.J., Geller R., Sparkes M.C. and Mohandas T. Regional mapping of human chromosome 19: Organisation of genes for plasma lipid transport (APOC1, -C2, and -E and LDLR) and the genes C3, PEPD and GPI. *Proc. Natl. Acad. Sci. USA.*, **83**, 3929-3933, 1986.
- Lynch P.J., Couch F.J., Gregg R.G., Powers P.A., Hogan K. and McCarthy T.V. Mutation screening of dihydropyridine receptor γ -subunit cDNA from malignant hyperthermia susceptible patients. *Biochem. Soc. Trans.*, **23**, 352S, 1995.

- Lynch P.J., Krivosic-Horber R., Reyford H., Monnier N., Quane K., Adnet P., Haudecoeur G., Krivosic I., McCarthy T. and Lunardi J. Identification of heterozygous and homozygous individuals with the novel RYR1 mutation Cys35Arg in a large kindred. *Anaesthesiology*, **86**, 620-626, 1997.
- Lynch P.J., Tong J., Lehane M., Mallet A., Giblin L., Heffron J.J.A., Vaughan P., Zafra G., MacLennan D.H. and McCarthy T.V. A mutation in the transmembrane/luminal domain of the ryanodine receptor is associated with abnormal Ca²⁺ release channel function and severe central core disease. *Proc. Natl. Acad. Sci. USA.*, **96**, 4164-4169, 1999.
- Mabry J.W., Christian L.L. and Kuhlert D.L. Inheritance of porcine stress syndrome. *J. Hered.*, **72**, 429-430, 1981.
- MacKenzie A.E., Korneluk R.G., Zorzato F., Fujii J., Phillips M., Iles D., Wieringa B., Leblond S., Bailly J., Willard H.F., Duff C., Worton R.G. and MacLennan D.H. The human ryanodine receptor gene: Its mapping to 19q13.1, placement in a chromosome 19 linkage group, and exclusion as the gene causing myotonic dystrophy. *Am. J. Hum. Genet.*, **46**, 1082-1089, 1990.
- Mackenzie A.E., Allen G., Lahey D., Crossan M-L., Nolan K., Mettler G., Worton R.G., MacLennan D.H. and Korneluk R. A comparison of the caffeine halothane muscle contracture test with the molecular genetic diagnosis of malignant hyperthermia. *Anesthesiology*, **75**, 4-8, 1991.
- MacLennan D.H. Purification and properties of an adenosine triphosphatase from sarcoplasmic reticulum. *J. Biol. Chem.*, **245**, 4508-4518, 1970.
- MacLennan D.H., Zorzato F., Fujii J., Otsu K., Phillips M., Lai F.A., Meissner G., Green N.M., Willard H.F., Britt B.A., Worton R.G. and Korneluk R.G. Cloning and localisation of the human calcium release channel (ryanodine receptor) gene to the proximal long arm (cen-q13.2) of human chromosome 19. *Am. J. Hum. Genet.*, **45**, A205, 1989.
- MacLennan D.H., Duff C., Zorzato F., Fujii J., Phillips M., Korneluk R.G., Frodis W., Britt B.A. and Worton R.G. Ryanodine receptor gene is a candidate for predisposition to malignant hyperthermia. *Nature*, **343**, 559-561, 1990.
- MacLennan D.H. and Phillips M.S. Malignant hyperthermia. *Science*, **256**, 789-794, 1992.
- Mader N., Gilly H. and Bittner R.E. Dystrophin deficient mdx muscle is not prone to MH susceptibility: An *in vitro* study. *Br. J. Anaesth.*, **79**, 125-127, 1997.
- Manning B.M., Quane K.A., Ording H., Urwyler A., Tegazzin V., Lehane M., O'Halloran J., Hartung E., Giblin L.M., Lynch P.J., Vaughan P., Censier K., Bendixen D., Comi G., Heytens L., Monsieurs K., Fagerlund T., Wolz W., Heffron J.J.A., Muller C.R. and McCarthy T.V. Identification of novel mutations in the ryanodine-receptor gene (RYR1) in malignant hyperthermia: Genotype-phenotype correlation. *Am. J. Hum. Genet.*, **62**, 599-609, 1998a.
- Manning B.M., Quane K.A., Lynch P.J., Urwyler A., Tegazzin V., Krivosic-Horber R., Censier K., Comi G., Abnet P., Wolz W., Lunardi J., Muller C.R. and McCarthy T.V. Novel mutations at a CpG dinucleotide in the ryanodine receptor in malignant hyperthermia. *Hum. Mutat.*, **11**, 45-50, 1998b.
- Marieb E.N. Human anatomy and physiology 3rd edition, The Benjamin/Cummings Publishing Company, Inc., Redwood City, California, USA., 1995.
- Martonosi A.N. and Pikula S. The structure of the Ca²⁺-ATPase of sarcoplasmic reticulum. *Acta Biochim. Pol.*, **50**, 337-365, 2003.
- Marx S.O., Gaburjakova J., Gaburjakova M., Henrikson C., Ondrias K. and Marks A.R. Coupled gating between cardiac calcium release channels (ryanodine receptors). *Circ. Res.*, **88**, 1151-1158, 2001.
- Masumiya H., Wang R., Zhang J., Xiao B. and Chen S.R. Localisation of the 12.6-kDa FK506-binding protein (FKBP12.6) binding site to the NH₂-terminal domain of the cardiac Ca²⁺ release channel (ryanodine receptor). *J. Biol. Chem.*, **278**, 3786-3792, 2003.
- McCarthy T.V., Healy J.M.S., Heffron J.J.A., Lehane M., Duefel T., Lehmann-Horn F., Farrall M. and Johnson K. Localisation of the malignant hyperthermia susceptibility locus to human chromosome 19q12-13.2. *Nature*, **343**, 562-564, 1990.
- McCarthy T.V., Quane K.A. and Lynch P.J. Ryanodine receptor mutations in malignant hyperthermia and central core disease. *Hum. Mutat.*, **15**, 410-417, 2000.
- McCarthy T.V. and Mackrill J.J. Unravelling calcium-release channel gating: Clues from a 'hot' disease. *Biochem. J.*, **380**, e1-e3, 2004.
- McPherson E.W. and Taylor C.A. The King syndrome: Malignant hyperthermia, myopathy and multiple anomalies. *Am. J. Med. Genet.*, **8**, 159-165, 1981.
- McPherson E. and Taylor C.A. The genetics of malignant hyperthermia: Evidence for heterogeneity. *Am. J. Med. Genet.*, **11**, 273-285, 1982.
- McPherson P.S. and Campbell K.P. The ryanodine receptor/Ca²⁺ release channel. *J. Biol. Chem.*, **268**, 13765-13768, 1993.
- McSweeney D.M. and Heffron J.J.A. Uptake and release of calcium ions by heavy sarcoplasmic reticulum fraction of normal and malignant hyperthermia-susceptible human skeletal muscle. *Int. J. Biochem.*, **22**, 329-333, 1990.
- McWilliams S., Nelson T., Sudo R.T., Zapata-Sudo G., Batti M. and Sambuughin N. Novel skeletal muscle ryanodine receptor mutation in a large Brazilian family with malignant hyperthermia. *Clin. Genet.*, **62**, 80-83, 2002.

- Meissner G. Ryanodine activation and inhibition of the Ca^{2+} release channel of sarcoplasmic reticulum. *J. Biol. Chem.*, **261**, 6300-6306, 1986.
- Menegazzi P., Larini F., Treves S., Guerrini R., Quadroni M. and Zorzato F. Identification and characterisation of three calmodulin binding sites of the skeletal muscle ryanodine receptor. *Biochemistry*, **33**, 9078-9084, 1994.
- Mezin P., Payen J-F., Bosson J-L., Brambilla E. and Stieglitz P. Histological support for the difference between malignant hyperthermia susceptible (MHS), equivocal (MHE) and negative (MHN) muscle biopsies. *Br. J. Anaesth.*, **79**, 327-331, 1997.
- Michalek-Sauberer A., Fricker R., Gradwohl I. and Gilly H. A case of suspected malignant hyperthermia during desflurane administration. *Anesth. Analg.*, **85**, 461-462, 1997.
- Mickelson J.R., Ross J.A., Reed B.K. and Louis C.F. Enhanced Ca^{2+} -induced calcium release by isolated sarcoplasmic reticulum vesicles from malignant hyperthermia susceptible pig muscle. *Biochim. Biophys. Acta.*, **862**, 318-328, 1986.
- Mickelson J.R., Gallant E.M., Litterer L.A., Johnson K.M., Rempel W.E. and Louis C.F. Abnormal sarcoplasmic reticulum ryanodine receptor in malignant hyperthermia. *J. Biol. Chem.*, **263**, 9310-9315, 1988.
- Mickelson J.R., Litterer L.A., Jacobson B.A. and Louis C.F. Stimulation and inhibition of [^3H]ryanodine binding to sarcoplasmic reticulum from malignant hyperthermia susceptible pigs. *Arch. Biochem. Biophys.*, **278**, 251-257, 1990.
- Mickelson J.R., Knudson C.M., Kennedy C.F.H., Yang D-I., Litterer L.A., Rempel W.E., Campbell K.P. and Louis C.F. Structural and functional correlates of a mutation in the malignant hyperthermia-susceptible pig ryanodine receptor. *FEBS. Lett.*, **301**, 49-52, 1992.
- Mieno S., Asada K., Horimoto H. and Sasaki S. Neuroleptic malignant syndrome following cardiac surgery: Successful treatment with dantrolene. *Eur. J. Cardiothorac. Surg.*, **24**, 458-460, 2003.
- Mitchell G. and Heffron J.J.A. Porcine stress syndromes. *Adv. Food Nutr. Res.*, **28**, 167-229, 1982.
- Monnier N., Procaccio V., Stieglitz P. and Lunardi J. Malignant hyperthermia susceptibility is associated with a mutation of the α_1 -subunit of the human dihydropyridine-sensitive L-type voltage-dependant calcium channel receptor in skeletal muscle. *Am. J. Hum. Genet.*, **60**, 1316-1325, 1997.
- Monnier N., Romero N.B., Lerale J., Nivoche Y., Qi D., MacLennan D.H., Fardeau M. and Lunardi J. An autosomal dominant congenital myopathy with cores and rods is associated with a neomutation in the RYR1 gene encoding the skeletal muscle ryanodine receptor. *Hum. Mol. Genet.*, **9**, 2599-2608, 2000.
- Monnier N., Romero N.B., Lerale J., Landrieu P., Nivoche Y., Fardeau M. and Lunardi J. Familial and sporadic forms of central core disease are associated with mutations in the C-terminal domain of the skeletal muscle ryanodine receptor. *Hum. Mol. Genet.*, **10**, 2581-2592, 2001.
- Monnier N., Krivosic-Horber R., Payen J-F., Kozak-Ribbens G., Nivoche Y., Adnet P., Reyford H. and Lunardi J. Presence of two different genetic traits in malignant hyperthermia families. *Anesthesiology*, **97**, 1067-1074, 2002.
- Monnier N., Ferreira A., Marty I., Labearre-Vila A., Mezin P. and Lunardi J. A homozygous splicing mutation causing a depletion of skeletal muscle RYR1 is associated with multi-minicore disease congenital myopathy with ophthalmoplegia. *Hum. Mol. Genet.*, **12**, 1171-1178, 2003.
- Monnier N., Kozak-Ribbens G., Krivosic-Horber R., Nivoche Y., Qi D., Kraev N., Loke J., Sharma P., Tegazzin V., Figarella-Branger D., Romero N., Mezin P., Bendahan D., Payen J.F., Depret T., MacLennan D.H. and Lunardi J. Correlations between genotype and pharmacological, histological, functional, and clinical phenotypes in malignant hyperthermia susceptibility. *Hum. Mutat.*, **26**, 413-425, 2005.
- Monsieurs K.G., Van Broeckhoven C., Martin J.J., Van Hoof V.O. and Heytens L. Gly341Arg mutation indicating malignant hyperthermia susceptibility: Specific cause of chronically elevated serum creatine kinase activity. *J. Neurol. Sci.*, **21**, 62-65, 1998.
- Moolman-Smook J.C., De Lange W.J., Bruwer E.C.D., Brink P.A. and Corfield V.A. The origins of hypertrophic cardiomyopathy – causing mutations in two South African subpopulations: A unique profile of both independent and founder events. *Am. J. Hum. Genet.*, **65**, 1308-1320, 1999.
- Moretti T., Koons B. and Budowle B. Enhancement of PCR amplification yield and specificity using AmpliTaq Gold™ DNA polymerase. *Biotechniques*, **25**, 716-722, 1998.
- Morgan P.G. and Sedensky M.M. Mutations causing a malignant hyperthermia-like state in *C.elegans*. *Anesthesiology*, **83**, 479, 1995.
- Moroni I., Gonano E.F., Comi G.P., Tegazzin V., Prella A., Bordoni A., Bresolin N. and Scarlato G. Ryanodine receptor gene point mutation and malignant hyperthermia susceptibility. *J. Neurol.*, **242**, 127-133, 1995.
- Moslehi R., Langlois S., Yam I. and Friedman J.M. Linkage of malignant hyperthermia and hyperkalemic periodic paralysis to the adult skeletal muscle sodium channel (SCN4A) gene in a large pedigree. *Am. J. Med. Genet.*, **76**, 21-27, 1998.

- Mouchiroud D., Gautier C. and Bernardi G. Frequencies of synonymous substitutions in mammals are gene-specific and correlated with frequencies of nonsynonymous substitutions. *J. Mol. Evol.*, **40**, 107-113, 1995.
- Mullis K., Faloona F., Scharf S., Saiki R., Horn G. and Erlich H. Specific enzymatic amplification of DNA *in vitro*: The polymerase chain reaction. *Cold Spring Harbour Symp. Quant. Biol.*, **L1**, 263-273, 1986.
- Murayama T. and Ogawa Y. Properties of RyR3 ryanodine receptor isoform in mammalian brain. *J. Biol. Chem.*, **271**, 5079-5084, 1996.
- Murayama T. and Ogawa Y. Characterisation of type 3 ryanodine receptor (RyR3) of sarcoplasmic reticulum from rabbit skeletal muscles. *J. Biol. Chem.*, **272**, 24030-24037, 1997.
- Nakai J., Imagawa T., Hakamat Y., Shigekawa M., Takeshima H. and Numa S. Primary structure and functional expression from cDNA of the cardiac ryanodine receptor/calcium release channel. *FEBS Lett.*, **27**, 169-177, 1990.
- Nakai J., Dirksen R.T., Nguyen H.T., Pessah I.N., Beam K.G. and Allen P.D. Enhanced dihydropyridine receptor channel activity in the presence of the ryanodine receptor. *Nature*, **380**, 72-75, 1996.
- Nakai J., Tanabe T., Konno T., Adams B. and Beam K.G. Localisation in the II-III loop of the dihydropyridine receptor of a sequence critical for excitation-contraction coupling. *J. Biol. Chem.*, **273**, 24983-24986, 1998.
- Nakayama H., Taki M., Striessnig J., Glossmann H., Catterall W.A. and Kanaoka Y. Identification of 1,4-dihydropyridine binding regions within the α_1 -subunit of skeletal muscle Ca^{2+} channels by photoaffinity labeling with diazepam. *Proc. Natl. Acad. Sci. USA.*, **88**, 9203-9207, 1991.
- Nelson T.E. and Flewelling E.H. Current concepts: The malignant hyperthermia syndrome. *N. Engl. J. Med.*, **18**, 416-418, 1983.
- Nelson T.E. Cell calcium in malignant hyperthermia. In: Cell calcium metabolism, physiology, biochemistry, pharmacology and clinical implications. Fiskum G., editor. Plenum Press, 507-511, 1989.
- Nelson T.E. Malignant hyperthermia in dogs. *J. Am. Vet. Med. A.*, **198**, 989-994, 1991.
- Nelson T.E., Lin M., Zapata-Sudo G. and Sudo R.T. Dantrolene sodium can increase or attenuate activity of skeletal muscle ryanodine receptor calcium release channel. Clinical implications. *Anesthesiology*, **84**, 1368-1379, 1996.
- Nelson T.E., Rosenberg H. and Muldoon S.M. Genetic testing for malignant hyperthermia in North America. *Anesthesiology*, **100**, 212-214, 2004.
- Neumann B.C. Molecular screening of the skeletal muscle ryanodine receptor (RYR1) gene for mutations associated with malignant hyperthermia susceptibility. M.Sc. Thesis, Potchefstroom University for Christian Higher Education, South Africa, 2002.
- Neylon C.B., Richards S.M., Larsen M.A., Agrotis A. and Bobik E. Multiple types of ryanodine receptor/ Ca^{2+} release channels are expressed in vascular smooth muscle. *Biochem. Biophys. Res. Commun.*, **215**, 814-821, 1995.
- Niebroj-Dobosz I. and Mayzner-Zawadzka E. Experimental porcine malignant hyperthermia: The activity of certain transporting enzymes and myofibrillar calcium-binding protein content in the muscle fibre. *Br. J. Anaesth.*, **54**, 885-890, 1982.
- Norborg M. and Tavaré S. Linkage disequilibrium: What history has to tell us. *Trends Genet.*, **18**, 83-90, 2002.
- O'Brien R.O., Take N.L., Hansbro P.M., Matthaai K.I., Hogan S.P., Denborough M.A. and Foster P.S. Exclusion of defects in the skeletal muscle specific regions of the DHPR α_1 -subunit as frequent cause of malignant hyperthermia. *J. Med. Genet.*, **32**, 913-914, 1995.
- O'Driscoll S., McCarthy T.V., Eichinger H.M., Erhardt W., Lehmann-Horn F. and Herrmann-Frank A. Calmodulin sensitivity of the sarcoplasmic reticulum ryanodine receptor from normal and malignant-hyperthermia-susceptible muscle. *Biochem. J.*, **319**, 421-426, 1996.
- Ohkoshi N., Yoshizawa T., Mizusawa H., Shoji S., Toyama M., Iida K., Sugishita Y., Hamano K., Takagi A., Goto K. and Arahata K. Malignant hyperthermia in a patient with Becker muscular dystrophy: Dytrophin analysis and caffeine contracture study. *Neuromusc. Disord.*, **5**, 53-58, 1995.
- Ohnishi S.T., Waring A.J., Fang S-R.G., Horiuchi K., Flick J.L., Sadanga K.K. and Ohnishi T. Abnormal membrane properties of the sarcoplasmic reticulum of pigs susceptible to malignant hyperthermia: Modes of action of halothane, caffeine, dantrolene and two other drugs. *Arch. Biochem. Biophys.*, **247**, 294-310, 1986.
- Ohnishi S.T., Katagi H., Ohnishi T. and Brownell A.K.W. Detection of malignant hyperthermia susceptibility using a spin label technique on red blood cells. *Br. J. Anaesth.*, **61**, 565-568, 1988.
- Oku S., Liew C-C. and Britt B.A. Analysis of sarcoplasmic reticulum proteins in patients susceptible to malignant hyperthermia. *J. Neurol. Sci.*, **60**, 127-135, 1983.
- Olckers A., Meyers D.A., Meyers S., Taylor E.W., Fletcher J.E., Rosenberg H., Isaacs H. and Levitt R.C. Adult sodium channel α -subunit is a gene candidate for malignant hyperthermia susceptibility. *Genomics*, **14**, 829-831, 1992.
- Olckers A. Molecular genetic analysis of the malignant hyperthermia susceptibility locus on human chromosome 19 in the South African malignant hyperthermia population. M.Sc. thesis, University of Pretoria, South Africa, 1993.

- Olickers A., Vita G.M., Rosenberg H., Fletcher J., Isaacs H. and Levitt R.C. Exclusion of the reported Arg⁶¹⁴ and Gly³⁴¹ ryanodine receptor proteins as being responsible for MHS in two families that map to chromosome 19q13.1, presented at the VIIth International Workshop on Malignant Hyperthermia, Hiroshima, Japan, 1994.
- Olickers A. Genetic heterogeneity in malignant hyperthermia susceptibility. Ph.D. thesis, University of Pretoria, South Africa, 1997.
- Olickers A., Meyers D.A., Taylor E., Havenga Y., Gericke G.S. and Weber J.L. Malignant hyperthermia: A possible novel locus on chromosome 2q suggested via linkage screening of the human genome, presented at the 8th Biennial Southern African Society of Human Genetics Congress, Gordon's Bay, South Africa, 1999.
- Olgin J., Argov Z., Rosenberg H., Tuchler M. and Chance B. Non-invasive evaluation of malignant hyperthermia susceptibility with phosphorus nuclear magnetic resonance spectroscopy. *Anesthesiology*, **68**, 507-513, 1988.
- Olgin J., Rosenberg H., Allen G., Seestedt R. and Chance B. A blinded comparison of noninvasive, *in vivo* phosphorus nuclear magnetic resonance spectroscopy and the *in vitro* halothane/caffeine contracture test in the evaluation of malignant hyperthermia susceptibility. *Anesth. Analg.*, **72**, 36-47, 1991.
- Ørding H. Incidence of malignant hyperthermia in Denmark. *Anesth. Analg.*, **64**, 700-704, 1985.
- Ørding H. and Bendixen D. Sources of variability in halothane and caffeine contracture tests for susceptibility to malignant hyperthermia. *Eur. J. Anaesthesiol.*, **9**, 367-376, 1992.
- Ørding H., Brancadoro V., Cozzolino S., Ellis F.R., Glauber V., Gonano E.F., Halsall P.J., Hartung E., Heffron J.J.A., Heytens L., Kozak-Ribbens G., Kress H., Krivosic-Horber R., Lehmann-Horn F., Mortier W., Nivoche Y., Ranklev-Twetman E., Sigurdsson S., Snoeck M., Stieglitz P., Tagazzin V., Urwyler A. and Wappler F. for the European Malignant Hyperthermia Group. *In vitro* contracture test for diagnosis of malignant hyperthermia following the protocol of the European MH Group: Results of testing patients surviving fulminant MH and unrelated low-risk subjects. *Acta. Anaesthesiol. Scand.*, **41**, 955-966, 1997.
- Otsu K., Nishida K., Kimura Y., Kuzuya T., Hori M., Kamada T. and Tada M. The point mutation Arg⁶¹⁵-Cys in the Ca²⁺ release channel of skeletal sarcoplasmic reticulum is responsible for hypersensitivity to caffeine and halothane in malignant hyperthermia. *J. Biol. Chem.*, **269**, 9413-9415, 1994.
- Ott J. Predicting the range of linkage disequilibrium. *P. Natl. Acad. Sci. USA.*, **97**, 2-3, 2000.
- Oyamada H., Oguchi K., Saitoh N., Yamazawa T., Hirose K., Kawana Y., Wakatsuki K., Oguchi K., Tagami M., Hanaoka K., Endo M. and Iino M. Novel mutations in C-terminal channel region of the ryanodine receptor in malignant hyperthermia patients. *Jpn. J. Pharmacol.*, **88**, 159-166, 2002.
- Paasuke R.T. and Brownell K.W. Serum creatine kinase level as a screening test for susceptibility to malignant hyperthermia. *J. Am. Med. Assoc.*, **255**, 769-771, 1986.
- Paavonen K.J., Chapman H., Laitinen P.J., Fodstad H., Piippo K., Swan H., Toivonen L., Viitasalo M., Kontula K. and Pasternack M. Functional characterisation of the common amino acid 897 polymorphism of the cardiac potassium channel KCNH2 (HERG). *Cardiovasc. Res.*, **59**, 603-611, 2003.
- Park J-W., Choi Y-B., Park S-K., Kim Y-I. and Lee K.S. Magnetic resonance imaging reveals selective vulnerability of the cerebellum and basal ganglia in malignant hyperthermia. *Arch. Neurol.*, **61**, 1462-1463, 2004.
- Parmley J.L., Chamary J.V. and Hurst L.D. Evidence for purifying selection against synonymous mutations in mammalian exonic splicing enhancers. *Mol. Biol. Evol.*, **23**, 301-309, 2005.
- Parness J. and Palnitkar S.S. Identification of dantrolene binding sites in porcine skeletal muscle sarcoplasmic reticulum. *J. Biol. Chem.*, **270**, 18465-18472, 1995.
- Patterson D.S.P. and Allen W.M. Biochemical aspects of some pig muscle disorders. *Br. Vet. J.*, **128**, 101-111, 1972.
- Paul-Pletzer K., Yamamoto T., Bhat M.B., Ma J., Ikemoto N., Jimenez L.S., Morimoto H., Williams P.G. and Parness J. Identification of the dantrolene binding sequence on the skeletal muscle ryanodine receptor. *J. Biol. Chem.*, **277**, 34918-34923, 2002.
- Perez C.F., Voss A., Pessah I.N. and Allen P.D. RyR1/RyR3 chimeras reveal that multiple domains of RyR1 are involved in skeletal-type E-C coupling. *Biophys. J.*, **84**, 2655-2663, 2003a.
- Perez C.F., Mukherjee S. and Allen P.D. Amino acids 1-1,680 of ryanodine receptor type 1 hold critical determinants of skeletal type for excitation-contraction coupling. Role of divergence domain D2. *J. Biol. Chem.*, **278**, 39644-39652, 2003b.
- Perez-Reyes E., Kim H.S., Lacerda A.E., Horne W., Wei X., Rampe D., Campbell K.P., Brown A.M. and Birnbaumer L. Induction of calcium currents by the expression of the α_1 -subunit of the dihydropyridine receptor from skeletal muscle. *Nature*, **340**, 233-236, 1989.
- Pessah I.N., Waterhouse A.L. and Casida J.E. The calcium-ryanodine receptor complex of skeletal and cardiac muscle. *Biochem. Biophys. Res. Commun.*, **128**, 449-456, 1985.
- Pessah I.N., Lynch C. and Gronert G.A. Complex pharmacology of malignant hyperthermia. *Anesthesiology*, **84**, 1275-1279, 1996.

- Petersen W.C. and Crouch E.R. Jr. Anesthesia-induced rigidity, unrelated to succinylcholine, associated with Smith-Lemli-Opitz syndrome and malignant hyperthermia. *Anesth. Analg.*, **80**, 606-608, 1995.
- Phillips M.S., Khanna V.K., De Leon S., Frodis W., Britt B.A. and MacLennan D.H. The substitution of Arg for Gly²⁴³³ in the human skeletal muscle ryanodine receptor is associated with malignant hyperthermia. *Hum. Mol. Genet.*, **3**, 2181-2186, 1994.
- Phillips M.S., Fujii J., Khanna V.K., DeLeon S., Yokobata K., De Jong P.J. and MacLennan D.H. The structural organisation of the human skeletal muscle ryanodine receptor (RYR1) gene. *Genomics*, **34**, 24-41, 1996.
- Pollock A.N., Langton E.E., Couchman K., Stowell K.M. and Waddington M. Suspected malignant hyperthermia reactions in New Zealand. *Anaesth. Intensive Care*, **30**, 453-461, 2002.
- Powers P.A., Scherer S.W., Tsui L-C., Gregg R.G. and Hogan K. Localisation of the gene encoding the α_2/δ -subunit (CACNL2A) of the human skeletal muscle voltage-dependent Ca^{2+} channel to chromosome 7q21-q22 by somatic cell hybrid analysis. *Genomics*, **19**, 192-193, 1994.
- Proenza C., Wilkens C., Lorenzon N.M. and Beam K.G. A carboxyl-terminal region important for the expression and targeting of the skeletal muscle dihydropyridine receptor. *J. Biol. Chem.*, **275**, 23169-23174, 2000.
- Proenza C., O'Brien J., Nakai J., Mukherjee S., Allen P.D. and Beam K.G. Identification of a region of RyR1 that participates in allosteric coupling with the α_{1S} ($Ca_v1.1$) II-III loop. *J. Biol. Chem.*, **227**, 6530-6535, 2002.
- QIAquick[®] Spin Handbook. Qiagen. QIAGEN Pty. Ltd., Victoria, Australia, 2002.
- Quane K.A., Healy J.M.S., Keating K.E., Manning B.M., Couch F.J., Palmucci L.M., Doriguzzi C., Fagerlund T.H., Berg K., Ordning H., Bendixen D., Mortier W., Linz U., Müller C.R. and McCarthy T.V. Mutations in the ryanodine receptor gene in central core disease and malignant hyperthermia. *Nat. Genet.*, **5**, 51-55, 1993.
- Quane K.A., Keating K.E., Manning B.M., Healy J.M.S., Monsieurs K., Heffron J.J.A., Lehane M., Heytens L., Krivosic-Horber R., Adnet P., Ellis F.R., Monnier N., Lunardi J. and McCarthy T.V. Detection of a novel common mutation in the ryanodine receptor gene in malignant hyperthermia: Implications for diagnosis and heterogeneity studies. *Hum. Mol. Genet.*, **3**, 471-476, 1994a.
- Quane K.A., Keating K.E., Healy J.M.S., Manning B.M., Krivosic-Horber R., Krivosic I., Monnier N., Lunardi J. and McCarthy T.V. Mutation screening of the RYR1 gene in malignant hyperthermia: Detection of a novel Tyr to Ser mutation in a pedigree with associated central cores. *Genomics*, **23**, 236-239, 1994b.
- Quane K.A., Ordning H., Keating K.E., Manning B.M., Heine R., Bendixen D., Berg K., Krivosic-Horber R., Lehmann-Horn F., Fagerlund T. and McCarthy T.V. Detection of a novel mutation at amino acid position 614 in the ryanodine receptor in malignant hyperthermia. *Br. J. Anaesth.*, **79**, 332-337, 1997.
- Rajabally Y.A. and El Lahawi M. Hypokalemic periodic paralysis associated with malignant hyperthermia. *Muscle Nerve*, **25**, 453-455, 2002.
- Rannala B. and Bertorelle G. Using linked markers to infer the age of a mutation. *Hum. Mutat.*, **18**, 87-100, 2001.
- Rasmussen U.F., Rasmussen H.N., Andersen A.J., Fogd Jørgensen P. and Quistorff B. Characterisation of mitochondria from pig muscle: Higher activity of exo-NADH oxidase in animals suffering from malignant hyperthermia. *Biochem. J.*, **315**, 659-663, 1996.
- Reed R. and Maniatis T. A role for exon sequences and splice-site proximity in splice-site selection. *Cell*, **46**, 681-690, 1986.
- Rees W.A., Yager T.D., Korte J. and Von Hippel P.H. Betaine can eliminate the base pair composition dependence of DNA melting. *Biochemistry*, **32**, 137-144, 1993.
- Richter M., Schleithoff L., Deufel T., Lehmann-Horn F. and Herrmann-Frank A. Functional characterisation of a distinct ryanodine receptor mutation in human malignant hyperthermia-susceptible muscle. *J. Biol. Chem.*, **272**, 5256-5260, 1997.
- Roberts V., Massimino M.L., Tosello V., Marsault R., Cantini M., Sorrentino V. and Pozzan T. Alteration in calcium handling at the subcellular level in *mdx* myotubes. *J. Biol. Chem.*, **276**, 4647-4651, 2001a.
- Roberts M.C., Mickelson J.R., Patterson E.E., Nelson T.E., Armstrong P.J., Brunson D.B. and Hogan K. Autosomal dominant canine malignant hyperthermia is caused by a mutation in the gene encoding the skeletal muscle calcium release channel (RYR1). *Anesthesiology*, **95**, 716-725, 2001b.
- Robinson R.L., Monnier N., Wolz W., Jung M., Reis A., Nuernberg G., Curran J.L., Monsieurs K., Stieglitz P., Heytens L., Fricker R., van Broeckhoven C., Deufel T., Hopkins P.M., Lunardi J. and Mueller C.R. A genome wide search for susceptibility loci in three European malignant hyperthermia pedigrees. *Hum. Mol. Genet.*, **6**, 953-961, 1997.
- Robinson R., Curran J.L., Hall W.J., Halsall P.J., Hopkins P.M., Markham A.F., Stewart A.D., West S.P. and Ellis F.R. Genetic heterogeneity and HOMOG analysis in British malignant hyperthermia families. *J. Med. Genet.*, **35**, 196-201, 1998.

- Robinson R.L., Curran J.L., Ellis F.R., Halsall P.J., Hall W.J., Hopkins P.M., Iles D.E., West S.P. and Shaw M-A. Multiple interacting gene products may influence susceptibility to malignant hyperthermia. *Ann. Hum. Genet.*, **64**, 307-320, 2000.
- Robinson R.L. and Hopkins P.M. A breakthrough in the genetic diagnosis of malignant hyperthermia. *Br. J. Anaesth.*, **86**, 166-168, 2001.
- Robinson R.L., Brooks C., Brown S.L., Ellis F.R., Halsall P.J., Quinnett R.J., Shaw M-A. and Hopkins P.M. RYR1 mutations causing central core disease are associated with more severe malignant hyperthermia *in vitro* contracture test phenotypes. *Hum. Mutat.*, **20**, 88-97, 2002.
- Robinson R.L., Anetseder M.J., Brancadoro V., van Broekhoven C., Carsana A., Censier K., Fortunato G., Girard T., Heytens L., Hopkins P.M., Jurkat-Rott K., Klinger W., Kozak-Ribbens G., Krivosic R., Monnier N., Nivoche Y., Olthoff D., Rueffert H., Sorrentino V., Tegazzin V. and Mueller C.R. Recent advances in the diagnosis of malignant hyperthermia susceptibility: How confident can we be of genetic testing? *Eur. J. Hum. Genet.*, **11**, 342-348, 2003a.
- Robinson R., Hopkins P., Carsana A., Gilly H., Halsall J., Heytens L., Islander G., Jurkat-Rott K., Müller C. and Shaw M-A. Several interacting genes influence the malignant hyperthermia phenotype. *Hum. Genet.*, **112**, 217-218, 2003b.
- Robinson R., Carpenter D., Shaw M-A., Halsall J. and Hopkins P. Mutations in RYR1 in malignant hyperthermia and central core disease. *Hum. Mutat.*, **27**, 977-989, 2006.
- Rogozin I.B., Spiridonov A.N., Sorokin A.V., Wolf Y.I., Jordan I.K., Tatusov R.L. and Koonin E.V. Purifying and directional selection in overlapping prokaryotic genes. *Trends Genet.*, **18**, 228-232, 2002.
- Romero N.B., Monnier N., Viollet L., Cortey A., Chevally M., Leroy J.P., Lunardi J. and Fardeau M. Dominant and recessive central core disease associated with RYR1 mutations and fetal akinesia. *Brain*, **126**, 2341-2349, 2003.
- Rosenberg H. and Fletcher J. International malignant hyperthermia workshop and symposium. *Anesthesiology*, **82**, 803-805, 1995.
- Rosenberg H. and Shutack J.G. Variants of malignant hyperthermia. Special problems for the paediatric anaesthesiologist. *Pediatr. Anaesth.*, **6**, 87-93, 1996.
- Rossi D. and Sorrentino V. The ryanodine receptors gene family: Expression and functional meaning. *Basic Appl. Myol.*, **14**, 323-343, 2004.
- Rueffert H., Olthoff D., Deutrich C., Kraus H. and Foster U.G. Identification of a novel polymorphism (IVS45+20 C/A) in the splice site of intron 45 of the ryanodine receptor gene (RYR1). *Hum. Mutat.*, Mutation and Polymorphism Report # **164**, 2000.
- Rueffert H., Olthoff D., Deutrich C., Thamm B. and Froster U.G. Homozygous and heterozygous Arg614Cys mutations (1840C-->T) in the ryanodine receptor gene co-segregate with malignant hyperthermia susceptibility in a German family. *Br. J. Anaesth.*, **87**, 240-245, 2001.
- Rueffert H., Olthoff D., Deutrich C., Meinecke C.D. and Froster U.G. Mutation screening in the ryanodine receptor 1 gene (RYR1) in patients susceptible to malignant hyperthermia who show definite IVCT results: Identification of three novel mutations. *Acta. Anaesthesiol. Scand.*, **46**, 692-698, 2002.
- Rueffert H., Olthoff D., Deutrich C., Schober R. and Froster U.G. A new mutation in the skeletal ryanodine receptor gene (RYR1) is potentially causative of malignant hyperthermia, central core disease, and severe skeletal malformation. *Am. J. Med. Genet.*, **30**, 248-54, 2004.
- Ruitenbeek W., Verburg M.P., Janssen A.J., Stadhouders A.M. and Sengers R.C. *In vivo* induced malignant hyperthermia in pigs. II. Metabolism of skeletal muscle mitochondria. *Acta. Anaesthesiol. Scand.*, **28**, 9-13, 1984.
- Ryan J.F. and Tedeschi L.G. Sudden unexplained death in a patient with a family history of malignant hyperthermia. *J. Clin. Anesth.*, **9**, 66-68, 1997.
- Sadusky T., Newman A.J. and Dibb N.J. Exon junction sequences as cryptic splice sites: Implications for intron origin. *Curr. Biol.*, **14**, 505-509, 2004.
- Saiki R.K., Gelfand D.H., Stoffel S., Scharf S.J., Higuchi R., Horn G.T., Mullis K.B. and Erlich H.A. Primer-directed enzymatic amplification of DNA with a thermostable DNA polymerase. *Science*, **239**, 487-491, 1988.
- Saito A., Inui M., Radermacher M., Frank J. and Fleischer S. Ultrastructure of the calcium release channel of sarcoplasmic reticulum. *J. Cell Biol.*, **107**, 211-219, 1988.
- Sambrook J.G. and Russell R. Molecular cloning: A laboratory manual. New York: Cold Spring Harbour Laboratory Press, Cold Spring Harbour, NY, USA, 2001.
- Sambuughin N., Sei Y., Gallagher K.L., Wyre H., Madsen D., Nelson T., Fletcher J., Rosenberg H. and Muldoon S. North American malignant hyperthermia population: Screening of the ryanodine receptor gene and identification of novel mutation. *Anesthesiology*, **95**, 594-599, 2001a.
- Sambuughin N., Nelson T.E., Jankovic J., Xin C., Meissner G., Mullakandov M., Ji J., Rosenberg H., Sivakumar K. and Goldfarb L.G. Identification and functional characterisation of a novel ryanodine receptor mutation causing malignant hyperthermia in North American and South American families. *Neuromusc. Dis.*, **11**, 530-537, 2001b.

- Sambuughin N., McWilliams S., de Bantel A., Sivakumar K. and Nelson T.E. Single-amino-acid deletion in the RYR1 gene, associated with malignant hyperthermia susceptibility and unusual contraction phenotype. *Am. J. Hum. Genet.*, **69**, 204-208, 2001c.
- Sambuughin N., Holley H., Muldoon S., Brandom B.W., de Bantel A.M., Tobin J.R., Nelson T.E. and Goldfarb L.G. Screening of the entire ryanodine receptor type 1 coding region for sequence variants associated with malignant hyperthermia susceptibility in the North American population. *Anesthesiology*, **102**, 515-521, 2005.
- Sanger F., Nicklen S. and Coulson A.R. DNA sequencing with chain-terminating inhibitors. *Proc. Natl. Acad. Sci. USA.*, **74**, 5463-5467, 1977.
- SantaLucia J. A unified view of polymer, dumbbell, and oligonucleotide DNA nearest-neighbour thermodynamics. *Proc. Natl. Acad. Sci. USA.*, **95**, 1460-1465, 1998.
- Sarker G., Kapelner S. and Sommer S.S. Formamide can dramatically improve the specificity of PCR. *Nucl. Acids Res.*, **18**, 7465, 1990.
- Saunders C.C. Historical dictionary of South Africa, African historical dictionaries no. 37. London, The Scarecrow Press xiii, 1983.
- Sawyer S.L., Mukherjee N., Pakstis A.J., Feuk L., Kidd J.R., Brookes A.J. and Kidd K.K. Linkage disequilibrium patterns vary substantially among populations. *Eur. J. Hum. Genet.*, **13**, 677-686, 2005.
- Scacheri P.C., Hoffman E.P., Fratkin J.D., Semino-Mora C., Senchak A., Davis M.R., Laing N.G., Vedanarayanan V. and Subramony S.H. A novel ryanodine receptor gene mutation causing both cores and rods in congenital myopathy. *Neurology*, **55**, 1689-1696, 2000.
- Schiller H.H. and Mair W.G.P. Ultrastructural changes of muscle in malignant hyperthermia. *J. Neurol. Sci.*, **21**, 93-100, 1974.
- Schönfeld P., Gerke S., Bohnensack R. and Wotjczak L. Stimulation of potassium cycling in mitochondrial by long-chain fatty acids. *Biochim. Biophys. Acta.*, **1604**, 125-133, 2003.
- Schübeler D. and Turner B.M. A new map for navigating the yeast epigenome. *Cell*, **122**, 489-492, 2005.
- Seay A.R. and Ziter F.A. Malignant hyperpyrexia in a patient with Schwartz-Jampel syndrome. *J. Pediatr.*, **93**, 83-84, 1978.
- Sedensky M.M. and Meneely P.M. Genetic analysis of halothane sensitivity in *Caenorhabditis elegans*. *Science*, **236**, 952-954, 1987.
- Sei Y., Gallagher K.L. and Basile A.S. Skeletal muscle type ryanodine receptor is involved in calcium signalling in human B lymphocytes. *J. Biol. Chem.*, **274**, 5995-6002, 1999.
- Sei Y., Sambuughin N.N., Davis E.J., Sachs D., Cuenca P.B., Brandom B.W., Tautz T., Rosenberg H., Nelson T.E. and Muldoon S.M. Malignant hyperthermia in North America. Genetic screening of the three hot spots in the type I ryanodine receptor gene. *Anesthesiology*, **101**, 824-830, 2004.
- Serfas K.D., Bose D., Patel L., Wrogemann K., Phillips M.S., MacLennan D.H. and Greenberg C.R. Comparison of the segregation of the RYR1 C1840T mutation with segregation of the caffeine/halothane contracture test results for malignant hyperthermia susceptibility in a large Manitoba Mennonite family. *Anesthesiology*, **84**, 322-329, 1996.
- Sewry C.A., Müller C., Davis M., Dwyer J.S.M., Dove J., Evans G., Schröder R., Fürst D., Helliwell T., Laing N. and Quinlivan R.C.M. The spectrum of pathology in central core disease. *Neuromuscul. Disord.*, **12**, 930-938, 2002.
- Sharma M.R. and Wagenknecht T. Cryo-electron microscopy and 3D reconstruction of ryanodine receptors and their interactions with E-C coupling proteins. *Basic Appl. Myol.*, **14**, 299-306, 2004.
- Shepherd S., Ellis F., Halsall J., Hopkins P. and Robinson R. RYR1 mutations in UK central core disease patients: More than just the C-terminal transmembrane region of the RYR1 gene. *J. Med. Genet.*, **41**, e33, 2004.
- Shoshan-Barmatz V., Orr I., Martin C. and Vardi N. Novel ryanodine-binding properties in mammalian retina. *Int. J. Biochem. Cell Biol.*, **37**, 1681-1695, 2005.
- Shy G.M. and Magee K.R. A new congenital non-progressive myopathy. *Brain*, **79**, 610-621, 1956.
- Smith J.S., Coronado R. and Meissner G. Single channel measurements of the calcium release channel from skeletal muscle sarcoplasmic reticulum. *J. Gen. Physiol.*, **88**, 573-588, 1986.
- Smith N.G.C., Webster M.T. and Ellegren H. Deterministic mutation rate variation in the human genome. *Genome Res.*, **12**, 1350-1356, 2002.
- Somers C.J. and McLoughlin J.V. Malignant hyperthermia in pigs: Calcium ion uptake by mitochondria from skeletal muscle of susceptible animals given neuroleptic drugs and halothane. *J. Comp. Path.*, **92**, 191-198, 1982.
- Sorrentino V., Giannini G., Malzac P. and Mattei M.G. Localisation of a novel ryanodine receptor gene (RYR3) to human chromosome 15q14-q15 by *in situ* hybridisation. *Genomics*, **18**, 163-165, 1993.
- Sorrentino V. and Reggiani C. Expression of the ryanodine receptor type 3 in skeletal muscle: A new partner in excitation-contraction coupling? *Trends Cardiovasc. Med.*, **9**, 54-61, 1999.
- Steinfath M., Singh S., Scholz J., Becker K., Lenzen C., Wappler F., Kochling A., Roewer N. and Schulte am Esch J. C1840-T mutation in the human skeletal muscle ryanodine receptor gene: Frequency in

- northern German families susceptible to malignant hyperthermia and the relationship to *in vitro* contracture response. *J. Mol. Med.*, **73**, 35-40, 1995.
- Steinmann C.M.L. A possible role of inositol 1,4,5-triphosphate (InsP₃) in malignant hyperthermia. *Med. Hypothesis*, **42**, 190-192, 1994.
- Stephan D.A. and Glueck S.B. Containing multitudes: Focus on "novel and nondetected human signaling protein polymorphisms". *Physiol. Genomics*, **10**, 127-129, 2002.
- Stewart C.R., Kahler S.G. and Gilchrist J.M. Congenital myopathy with cleft palate and increase susceptibility to malignant hyperthermia: King syndrome? *Pediatr. Neurol.*, **4**, 371-374, 1988.
- Stewart S.L., Rosenberg H. and Fletcher J.E. Failure to identify the ryanodine receptor G1021A mutation in a large North American population with malignant hyperthermia. *Clin. Genet.*, **54**, 358-361, 1998.
- Stewart S.L., Hogan K., Rosenberg H. and Fletcher J.E. Identification of the Arg1086His mutation in the alpha subunit of the voltage-dependant calcium channel (CACNA1S) in a North American family with malignant hyperthermia. *Clin. Genet.*, **59**, 178-184, 2001.
- Strazis K.P. and Fox A.W. Malignant hyperthermia: A review of published cases. *Anesth. Analg.*, **77**, 297-304, 1993.
- Sudbrak R., Procaccio V., Klausnitzer M., Curran J.L., Monsieus K., Van Broeckhoven C., Ellis R., Heyetens L., Hartung E.J., Kozak-Ribbens G., Heilinger D., Weissenbach J., Lehman-Horn F., Mueller C.R., Deufel T., Stewart A.D. and Lunardi J. Mapping of a further malignant hyperthermia susceptibility locus to chromosomes 3q13.1. *Am. J. Hum. Genet.*, **56**, 684-691, 1995.
- Sugimoto N., Nakano S., Yoneyama M. and Honda K. Improved thermodynamic parameters and helix initiation factor to predict stability of DNA duplexes. *Nucleic Acids Res.*, **24**, 4501-4505, 1996.
- Suko J., Maurer-Fogy I., Plank B., Bertel O., Wyskovsky W., Hohenegger M. and Hellmann G. Phosphorylation of serine 2843 in ryanodine receptor-calcium release channel of skeletal muscle by cAMP-, cGMP- and CaM-dependant protein kinase. *Biochim. Biophys. Acta.*, **1175**, 193-206, 1993.
- Sun J., Xin C., Eu J.P., Stamler J.S. and Meissner G. Cysteine-3635 is responsible for skeletal muscle ryanodine receptor modulation by NO. *Proc. Natl. Acad. Sci. USA.*, **98**, 11158-11162, 2001.
- Suresh M.S. and Nelson T.E. Malignant hyperthermia: Is etomidate safe? *Anesth. Analg.*, **64**, 420-424, 1985.
- Takehima H., Nishimura S., Matsumoto T., Ishida H., Kangawa K., Minamino N., Matsuo H., Ueda M., Hanaoka M., Hirose T. and Numa S. Primary structure and expression from complementary DNA of skeletal muscle ryanodine receptor. *Nature*, **339**, 439-445, 1989.
- Tamaro A., Bracco A., Cozzolino S., Esposito M., Di Matino A., Savoia G., Zeuli L., Piluso G., Aurino S. and Nigro V. Scanning for mutations of the ryanodine receptor (RYR1) gene by denaturing HPLC: Detection of three novel malignant hyperthermia alleles. *Clin. Chem.*, **49**, 761-768, 2003.
- Tanabe T., Beam K.G., Adams B.A., Niidome T. and Numa S. Regions of the skeletal muscle dihydropyridine receptor critical for excitation-contraction coupling. *Nature*, **346**, 567-469, 1990.
- Tilgen N., Zorzato F., Halliger-Keller B., Muntoni F., Sewry C., Palmucci L.M., Schneider C., Hauser E., Lehmann-Horn F., Müller C.R. and Treves S. Identification of four novel mutations in the C-terminal membrane spanning domain of the ryanodine receptor 1: Association with central core disease and alteration of calcium homeostasis. *Hum. Mol. Genet.*, **10**, 2879-2887, 2001.
- Timerman A.P., Ogunbumni E., Freund E., Wiederrecht G., Marks A.R. and Fleischer S. The calcium release channel of sarcoplasmic reticulum is modulated by FK-506-binding protein. *J. Biol. Chem.*, **268**, 22992-22999, 1993.
- Tipping A.J., Pearson T., Morgan N.V., Gibson R.A., Kuyt L.P., Havenga C., Gluckman E., Joenje H., de Ravel T., Jansen S. and Mathew C.G. Molecular and genealogical evidence for a founder effect in Faconi anemia families of the Afrikaner population in South Africa. *P. Natl. Acad. Sci. USA.*, **98**, 5734-5739, 2001.
- Tiso N., Stephan D.A., Nava A., Bagattin A., Devaney J.M., Stanchi F., Larderet G., Brahmabhatt B., Brown K., Bauce B., Muriago M., Basso C., Thiene G., Danieli G.A. and Rampazzo A. Identification of mutations in the cardiac ryanodine receptor gene in families affected with arrhythmogenic right ventricular cardiomyopathy type 2 (ARVD2). *Hum. Mol. Genet.*, **10**, 189-194, 2001.
- Tong J., Oyamada H., Demaurex N., Grinstein S., McCarthy T.V. and MacLennan D.H. Caffeine and halothane sensitivity of intracellular Ca²⁺ release is altered by 15 calcium release channel (ryanodine receptor) mutations associated with malignant hyperthermia and/or central core disease. *J. Biol. Chem.*, **272**, 26332-26339, 1997.
- Tonner P.H., Scholz J., Richter A., Löscher W., Steinfath M., Wappler F., Wlaz P., Hadji B., Roewer N. and Schulte am Esch J. Alterations of inositol polyphosphates in skeletal muscle during porcine malignant hyperthermia. *Br. J. Anaesth.*, **75**, 467-471, 1995.
- Treves S., Larini F., Menegazzi P., Steinberg T.H., Koval M., Vilsen B., Andersen J.P. and Zorzato F. Alteration of intracellular Ca²⁺ transients in COS-7 cells transfected with the cDNA encoding skeletal-muscle ryanodine receptor carrying a mutation associated with malignant hyperthermia. *Biochem J.*, **301**, 661-665, 1994.

- Treves S., Pouliquin P., Moccagatta L. and Zorzato F. Functional properties of EGFP-tagged skeletal muscle calcium-release channel (ryanodine receptor) expressed in COS-7 cells: Sensitivity to caffeine and 4-chloro-*m*-cresol. *Cell Calcium*, **31**, 1-12, 2002.
- Tunwell R.E.A., Wickenden C., Bertrand B.M.A., Shevchenko V.I., Walsh M.B., Allen P.D. and Lai F.A. The human cardiac muscle ryanodine receptor-calcium release channel: Identification, primary structure and topological analysis. *Biochem. J.*, **318**, 477-487, 1996.
- Urwyler A., Deufel T., McCarthy T. and West S. for the European Malignant Hyperthermia Group. Guidelines for molecular genetic detection of susceptibility to malignant hyperthermia. *Br. J. Anaesth.*, **86**, 283-287, 2001.
- Valdivia H.H., Hogan K. and Cornado R. Altered binding site for Ca²⁺ in the ryanodine receptor of human malignant hyperthermia. *Am. J. Physiol.*, **261**, C237-C245, 1991.
- Verona R.I., Mann M.R.W. and Bartolmei M.S. Genomic imprinting: Intricacies of epigenetic regulation in clusters. *Ann. Rev. Cell Dev. Biol.*, **19**, 237-259, 2003.
- Vita G.M., Olckers A., Jedlicka A.E., George A.L., Heiman-Patterson T., Rosenberg H., Fletcher J.E. and Levitt R.C. Masseter muscle rigidity associated with Glycine¹³⁰⁶-to-Alanine mutation in the adult muscle sodium channel α -subunit gene. *Anesthesiology*, **82**, 1097-1103, 1995.
- Voet D. and Voet J.G. Biochemistry 2nd edition, John Wiley and Sons, Somerset, New Jersey, USA., 1999.
- Voss A.A., Lango J., Ernst-Russell M., Morin D. and Pessah I.N. Identification of hyperreactive cysteines within ryanodine receptor type 1 by mass spectroscopy. *J. Biol. Chem.*, **279**, 34514-34520, 2004.
- Wagenknecht T., Grassucci R., Frank J., Saito A., Inui M. and Fleischer S. Three-dimensional architecture of the calcium channel/foot structure of sarcoplasmic reticulum. *Nature*, **338**, 167-170, 1989.
- Wagenknecht T. and Radermacher M. Three-dimensional architecture of the skeletal muscle ryanodine receptor. *FEBS Lett.*, **369**, 43-46, 1995.
- Wagenknecht T., Radermacher M., Grassucci R., Berkowitz J., Xin H-B. and Fleischer S. Locations of calmodulin and FK506-binding protein on the three-dimensional architecture of the skeletal muscle ryanodine receptor. *J. Biol. Chem.*, **272**, 32436-32471, 1997.
- Wang D. and Sadée W. Searching for polymorphisms that affect gene expression and mRNA processing: Example ABCB1 (MDR1). *AAPS J.*, **8**, 515-520, 2006.
- Wappler F., Roewer N., Köchling A., Braune H., Reissinger T. and Schulte am Esch J. Fulminant malignant hyperthermia associated with ketoacidotic diabetic coma. *Intensive Care Med.*, **22**, 809-812, 1996.
- Wappler F. Malignant hyperthermia. *Eur. J. Anaesth.*, **18**, 632-652, 2001.
- Wedel D.J., Gammel S.A., Milde J.H. and Iazzo P.A. Delayed onset of malignant hyperthermia induced by isoflurane and desflurane compared with halothane in susceptible swine. *Anesthesiology*, **78**, 1138-1144, 1993.
- Wehner M., Rueffert H., Koenig F., Neuhaus J. and Olthoff D. Increased sensitivity to 4-chloro-*m*-cresol and caffeine in primary myotubes from malignant hyperthermia susceptible individuals carrying the ryanodine receptor 1 Thr2206Met (C6617T) mutation. *Clin. Genet.*, **62**, 135-146, 2002.
- Wehner M., Rueffert H., Koenig F., Meinecke C.D. and Olthoff D. The Ile2453Thr mutation in the ryanodine receptor gene 1 is associated with facilitated calcium release from sarcoplasmic reticulum by 4-chloro-*m*-cresol in human myotubes. *Cell Calcium*, **34**, 163-168, 2003.
- Wehner M., Rueffert H., Koenig F. and Olthoff D. Functional characterisation of malignant hyperthermia-associated RyR1 mutations in exon 44, using the human myotube model. *Neuromuscul. Disord.*, **14**, 429-437, 2004.
- Weiss R.G., O'Connell K.M., Flucher B.E., Allen P.D., Grabner M. and Dirksen R.T. Functional analysis of the R1086H malignant hyperthermia mutation in the DHPR reveals an unexpected influence of the III-IV loop on skeletal muscle EC coupling. *Am. J. Physiol. Cell. Physiol.*, **287**, 1094-1102, 2004.
- Wingard D.W. Malignant hyperthermia: A human stress syndrome? *Lancet*, **2**, 1450-1451, 1974.
- Wittwer C.T., Reed G.H., Gundry C.N., Vandersteen J.G. and Pryor R.J. High-resolution genotyping by amplicon melting analysis using LCGreen. *Clin. Chem.*, **49**, 853-860, 2003.
- Wizard[®] Genomic DNA Purification Kit Technical Manual. Promega Corporation, Madison, USA, 1998.
- Wrogemann K. and Pena S.D.J. Mitochondrial calcium overload: A general mechanism for cell-necrosis in muscle diseases. *Lancet*, **1**, 672-674, 1976.
- Wu D.Y., Ugozzoli L., Pal B.K., Qian J. and Wallace R.B. The effect of temperature and oligonucleotide primer length on the specificity and efficiency of amplification by the polymerase chain reaction. *DNA Cell Biol.*, **10**, 233-238, 1991.
- Wu S., Ibarra C.A., Malicdan M.C.V., Murayama K., Ichihara Y., Kikuchi H., Nonaka I., Noguchi S., Hayashi Y.K. and Nishino I. Central core disease is due to RYR1 mutations in more than 90% of patients. *Brain*, **129**, 1470-1480, 2006.
- Yamamoto T., El-Hayek R. and Ikemoto N. Postulated role of interdomain interaction within the ryanodine receptor in Ca²⁺ channel regulation. *J. Biol. Chem.*, **275**, 11618-11625, 2000.
- Yang T., Ta T.A., Pessah I.N. and Allen P.D. Functional defects in six ryanodine receptor isoforms-1 (RyR1) mutations associated with malignant hyperthermia and their impact on skeletal excitation-contraction coupling. *J. Biol. Chem.*, **278**, 25722-25730, 2003.

- Yaqub B. and Al Deeb S. Heat strokes: Aetiopathogenesis, neurological characteristics, treatment and outcome. *J. Neurolog. Sci.*, **156**, 144-151, 1998.
- Yeh H.M., Tsai M.C., Su Y.N., Shen R.C., Hwang J.J., Sun W.Z. and Lai L.P. Denaturing high performance liquid chromatography screening of the ryanodine receptor type 1 gene in patients with malignant hyperthermia in Taiwan and identification of a novel mutation (Y522C). *Anesth. Analg.*, **101**, 1401-1406, 2005.
- Zhang Y., Chen H.S., Khanna V.K., De Leon S., Phillips M.S., Schappert K., Britt B.A., Brownell K.W. and MacLennan D.H. A mutation in the human ryanodine receptor gene associated with central core disease. *Nat. Genet.*, **5**, 46-50, 1993.
- Zhang L., Kelley J., Schmeisser G., Kobayashi Y.M. and Jones L.R. Complex formation between junctin, triadin, calsequestrin, and the ryanodine receptor. *J. Biol. Chem.*, **272**, 23389-23397, 1997.
- Zhou H., Treves S., Jungbluth H., Sewry C. and Mutoni F. RYR1 gene genotype-phenotype and functional correlative studies in central core disease and multi-minicore disease. *Neuromuscul. Disord.*, **15**, 676-743, 2005.
- Zhou H., Yamaguchi N., Xu L., Wang Y., Sewry C., Jungbluth H., Zorzato F., Bertini E., Muntoni F., Meissner G. and Treves S. Characterisation of recessive RYR1 mutations in core myopathies. *Hum Mol Genet.*, **15**, 2791-2803, 2006a.
- Zhou H., Brockington M., Jungbluth H., Monk D., Stanier P., Sewry C.A., Moore G.E. and Muntoni F. Epigenetic allele silencing unveils recessive RYR1 mutations in core myopathies. *Am. J Hum. Genet.*, **79**, 859-868, 2006b.
- Zhu X., Ghanta J., Walker J.W., Allen P.D. and Valdivia H.H. The calmodulin binding region of the skeletal ryanodine receptor acts as a self-modulatory domain. *Cell Calcium*, **35**, 165-177, 2004.
- Zorzato F., Fujii J., Otsu K., Phillips M., Green N.M., Lai F.A., Meissner G. and MacLennan D.H. Molecular cloning of cDNA encoding human and rabbit forms of the Ca²⁺ release channel (ryanodine receptor) of skeletal muscle sarcoplasmic reticulum. *J. Biol. Chem.*, **265**, 2244-2256, 1990.
- Zorzato F., Yamaguchi N., Xu L., Meissner G., Müller C.R., Pouliquin P., Muntoni F., Sewry C., Girard T., and Treves S. Clinical and functional effects of a deletion in a COOH-terminal luminal loop of the skeletal muscle ryanodine receptor. *Hum. Mol. Genet.*, **12**, 379-388, 2003.
- Zuker M. Mfold web server for nucleic acid folding and hybridization prediction. *Nucleic Acids Res.*, **31**, 3406-3415, 2003.

6.2 ELECTRONIC REFERENCES

- BioEdit, Biological sequence alignment editor, 9 June 2005, <http://www.mbio.ncsu.edu/Bioedit/biedit/html>
- European Molecular Biology Laboratory, 9 November 2005, <http://www.ebi.ac.uk/>
- GENBANK, 15 July 2004, <http://ncbi.nlm.nih.gov/Genbank/>
- Harbitz I., Kristensen T., Bosnes M., Kran S. and Davies W. DNA sequence of the skeletal muscle calcium release channel cDNA and verification of the Arg615-Cys615 mutation, associated with porcine malignant hyperthermia, in Norwegian landrace pigs. GenBank accession number P16960, 1992.
- Integrated DNA Technologies, 10 November 2005, <http://www.idtdna.com/SciTools/SciTools.aspx>
- International Human Genome Sequencing Consortium. Finishing the euchromatic sequence of the human genome. *Nature*, **431**, 931-945, 2004, with accession number rs1465698, 10 November 2005, <http://ncbi.nlm.nih.gov/Genbank/>
- International Human Genome Sequencing Consortium. Finishing the euchromatic sequence of the human genome. *Nature*, **431**, 931-945, 2004, with accession number rs1465698, 10 November 2005, <http://ncbi.nlm.nih.gov/Genbank/>
- International Human Genome Sequencing Consortium. Finishing the euchromatic sequence of the human genome. *Nature*, **431**, 931-945, 2004, with accession number rs2071085, 10 November 2005, <http://ncbi.nlm.nih.gov/Genbank/>
- International Human Genome Sequencing Consortium. Finishing the euchromatic sequence of the human genome. *Nature*, **431**, 931-945, 2004, with accession number rs2071088, 10 November 2005, <http://ncbi.nlm.nih.gov/Genbank/>
- International Human Genome Sequencing Consortium. Finishing the euchromatic sequence of the human genome. *Nature*, **431**, 931-945, 2004, with accession number rs2071089, 10 November 2005, <http://ncbi.nlm.nih.gov/Genbank/>
- International Human Genome Sequencing Consortium. Finishing the euchromatic sequence of the human genome. *Nature*, **431**, 931-945, 2004, with accession number rs2228068, 10 November 2005, <http://ncbi.nlm.nih.gov/Genbank/>

- International Human Genome Sequencing Consortium. Finishing the euchromatic sequence of the human genome. *Nature*, **431**, 931-945, 2004, with accession number rs10406027, 10 November 2005, <http://ncbi.nlm.nih.gov/Genbank/>
- International Human Genome Sequencing Consortium. Finishing the euchromatic sequence of the human genome. *Nature*, **431**, 931-945, 2004, with accession number rs11083462, 10 November 2005, <http://ncbi.nlm.nih.gov/Genbank/>
- International Human Genome Sequencing Consortium. Finishing the euchromatic sequence of the human genome. *Nature*, **431**, 931-945, 2004, with accession number rs12460768, 10 November 2005, <http://ncbi.nlm.nih.gov/Genbank/>
- International Human Genome Sequencing Consortium. Finishing the euchromatic sequence of the human genome. *Nature*, **431**, 931-945, 2004, with accession number rs12461853, 10 November 2005, <http://ncbi.nlm.nih.gov/Genbank/>
- International Human Genome Sequencing Consortium. Finishing the euchromatic sequence of the human genome. *Nature*, **431**, 931-945, 2004, with accession number rs12973632, 10 November 2005, <http://ncbi.nlm.nih.gov/Genbank/>
- International Human Genome Sequencing Consortium. Finishing the euchromatic sequence of the human genome. *Nature*, **431**, 931-945, 2004, with accession number rs12985668, 10 November 2005, <http://ncbi.nlm.nih.gov/Genbank/>
- International Human Genome Sequencing Consortium. Finishing the euchromatic sequence of the human genome. *Nature*, **431**, 931-945, 2004, with accession number rs17708009, 10 November 2005, <http://ncbi.nlm.nih.gov/Genbank/>
- International Human Genome Sequencing Consortium. Finishing the euchromatic sequence of the human genome. *Nature*, **431**, 931-945, 2004, with accession number rs35959206, 10 November 2005, <http://ncbi.nlm.nih.gov/Genbank/>
- Malignant Hyperthermia Association of the United States, 9 November 2005, <http://www.mhaus.org/index.cfm/fuseaction/Content.Display/PagePK/Home.cfm>
- Oligonucleotide properties calculator, 21 May 2005, <http://www.basic.nwu.edu/biotools/oligocalc.html>, Xprimer, Department of Genetics and Cell Biology, University of Minnesota, <http://www.alces.med.umn.edu/rawprimers>
- Takeshima H., Nishimura S., Matsumoto T., Ishido H., Kangawa K., Minamino N., Matsuo H., Ueda M., Hanaoka M., Hirose T. and Numa S. Primary structure and expression from complementary DNA of skeletal muscle ryanodine receptor. GenBank accession number P11716, 1989.
- Tunwell R.E., Wickenden C., Bertrand B.M., Shevchenko V.I., Walsh M.B., Allen P.D. and Lai F.A. The human cardiac muscle ryanodine receptor-calcium release channel: Identification, primary structure and topological analysis. GenBank accession number Q92736, 1996.
- Van Acker K., Bultynck G., Rossi D., Sorrentino V., Boens N., Missiaen L., De Smedt H., Parys J.B. and Callewaert G. The 12 kDa FK506-binding protein, FKBP12, modulates the Ca²⁺-flux properties of the type-3 ryanodine receptor. GenBank accession number NP_001027, 2004.
- Zorzato F., Fujii J., Otsu K., Phillips M., Green N.M., Lai F.A., Meissner G. and MacLennan D.H. Molecular cloning of cDNA encoding human and rabbit forms of the Ca²⁺ release channel (ryanodine receptor) of skeletal muscle sarcoplasmic reticulum. GenBank accession number P21817, 1990.

