

Diversity of fall armyworm, *Spodoptera frugiperda* and their gut bacterial community in Kenya

Joseph Gichuhi¹, Subramanian Sevgan¹, Fathiya Khamis¹,
Johnnie Van den Berg², Hannalene du Plessis², Sunday Ekesi¹ and
Jeremy K. Herren^{1,3}

¹ International Centre of Insect Physiology and Ecology (ICIPE), Nairobi, Kenya

² Unit for Environmental Sciences and Management, North-West University, Potchefstroom, South Africa

³ MRC-University of Glasgow Centre for Virus Research, Henry Wellcome Building, Glasgow, UK

ABSTRACT

Background: The invasive fall armyworm, *Spodoptera frugiperda* (J.E. Smith) is a polyphagous pest that causes widespread damage particularly to maize and sorghum in Africa. The microbiome associated with *S. frugiperda* could play a role in the insects' success and adaptability. However, bacterial communities in *S. frugiperda* remain poorly studied.

Methods: We investigated the composition, abundance and diversity of microbiomes associated with larval and adult specimens of *S. frugiperda* collected from four maize growing regions in Kenya through high throughput sequencing of the bacterial 16S rRNA gene. The population structure of *S. frugiperda* in Kenya was assessed through amplification of the mitochondrial cytochrome oxidase subunit I gene.

Results: We identified Proteobacteria and Firmicutes as the most dominant bacterial phyla and lesser proportions of Bacteroidetes and Actinobacteria. We also observed differences in bacterial microbiome diversity between larvae and adults that are a likely indication that some prominent larval bacterial groups are lost during metamorphosis. However, several bacterial groups were found in both adults and larvae suggesting that they are transmitted across developmental stages. Reads corresponding to several known entomopathogenic bacterial clades as well as the fungal entomopathogen, *Metarhizium rileyi*, were observed. Mitochondrial DNA haplotyping of the *S. frugiperda* population in Kenya indicated the presence of both "Rice" and "Corn" strains, with a higher prevalence of the "Rice" strain.

Subjects Agricultural Science, Entomology, Microbiology, Molecular Biology

Keywords Fall armyworm, *Spodoptera frugiperda*, Gut bacteria, Corn strain, Rice strain, mtCOI haplotype

INTRODUCTION

Invasions by exotic pests can have major detrimental effects on agricultural production and natural resources (Huber *et al.*, 2002). The fall armyworm (FAW), *Spodoptera frugiperda* (J.E. Smith) (Lepidoptera: Noctuidae) is a polyphagous pest that is, native to tropical regions of the western hemisphere, where it is known for its ability to cause economic damage to several crop species. In 2016, *S. frugiperda* was first detected in

Submitted 21 October 2019

Accepted 6 February 2020

Published 5 March 2020

Corresponding author

Jeremy K. Herren, jherren@icipe.org

Academic editor

Brenda Oppert

Additional Information and
Declarations can be found on
page 14

DOI 10.7717/peerj.8701

© Copyright

2020 Gichuhi et al.

Distributed under

Creative Commons CC-BY 4.0

OPEN ACCESS

West Africa (Goergen *et al.*, 2016), and since then this pest has rapidly spread across the continent (Day *et al.*, 2017; Nagoshi *et al.*, 2018; Rwomushana *et al.*, 2018). By 2018, *S. frugiperda* was reported in all countries in Sub-Saharan Africa except Djibouti and Lesotho (Rwomushana *et al.*, 2018). Furthermore, *S. frugiperda* also has now reached the continent of Asia (Deole & Paul, 2018; Sisodiya *et al.*, 2018). Maize and other economically important food crops in these regions are extensively damaged by *S. frugiperda* larvae (Day *et al.*, 2017) causing extensive economic losses and threatening food security. Genetic characterizations have supported the initial postulation that this pest species exists in two subpopulations called the “Rice” and “Corn” strains that are preferentially associated either with smaller grasses (such as rice and bermudagrass) or with larger grasses (such as sorghum and maize) respectively (Nagoshi & Meagher, 2004). Strain composition in populations of this pest may therefore have ramifications on the variety of crops at risk of infestation (Nagoshi *et al.*, 2019).

There is a lack of information about *S. frugiperda*-host plant interactions and other factors that may be leading to the rapid spread of *S. frugiperda* in the geographic regions that have recently been invaded. Many of the control measures used in the western hemisphere (e.g., transgenic maize, chemical insecticides) might not be readily available and economically viable for subsistence farmers in Africa. Furthermore, the use of highly hazardous pesticides is not considered a sustainable long term control measure for any pest (FAO, 2018). In addition, *S. frugiperda* have been reported to evolve resistance to most chemical insecticides (e.g., pyrethroids, organophosphates and carbamates) (Yu, 1991) and to transgenic maize that are used in its control (Jakka *et al.*, 2016; Banerjee *et al.*, 2017; Flagel *et al.*, 2018; Botha *et al.*, 2019). As a consequence, there is a great need for alternative, cost-effective control strategies for *S. frugiperda* (FAO, 2018).

A recent survey in Ethiopia, Kenya and Tanzania indicated that *S. frugiperda* has established interactions with indigenous parasitoid species (Sisay *et al.*, 2018) that could be harnessed for biological control. A study on *S. frugiperda* host plant interactions in East Africa has also suggested a climate adapted push-pull system (Midtega *et al.*, 2018) and maize-legume intercropping (Hailu *et al.*, 2018) for management of pests including FAW on maize farms. However, many factors related to *S. frugiperda* rapid spread, host plant interactions, bio-ecology and insect-microbiome interactions in the African region remain poorly understood.

Insect microbiomes can have important consequences for the outcome of insect pest-natural enemies- host plant interactions (Ferrari, Vavre & Lyon, 2011). Strategies that involve modifying insect microbiomes are currently being evaluated for control and management of pests and vectors of plant diseases (Crotti *et al.*, 2012; Perilla-Henao & Casteel, 2016; Arora & Douglas, 2017; Beck & Vannette, 2017). Insect microbiomes play a key role in the adaptation of insects to their environment and are therefore a major and often poorly understood determinant of the host plant and geographic range of insect pests (Su, Zhou & Zhang, 2013). In general, a greater diversity of microbial symbionts exist within the insect’s gut lumen, while few others exist inside cells of the host, or on the cuticle (Douglas, 2015). Gut microbial symbionts are known to influence their host’s nutrition, usually by promoting digestion and availability of nutrients (Douglas, 2009).

These symbionts can also modulate the immune response and accessibility of the host to invading organisms, and therefore have direct or indirect effects on host susceptibility to parasites and pathogens ([Garcia et al., 2010](#); [McLean & Godfray, 2015](#); [Ubeda, Djukovic & Isaac, 2017](#)). Previous studies have also identified important roles of bacterial symbionts in the interactions between phytophagous insects and host plants ([Frago, Dicke & Godfray, 2012](#); [Biere & Bennett, 2013](#); [Brady & White, 2013](#)). In addition, microbial symbionts can break down complex molecules such as insecticides and promote insecticide resistance ([Kikuchi et al., 2012](#); [Xia et al., 2018](#)). It is also notable that pathogenic bacteria can reside in host guts, only initiating or facilitating pathogenesis under certain conditions ([Wei et al., 2017](#)). Studying the gut microbiome is not only important from the standpoint of understanding mutualistic relationships but also for laying the foundation for future projects aimed at developing microbial biocontrol agents.

There are an increasing number of studies examining the microbial diversity of lepidopterans. While in some of the assessed species consistent bacterial communities have been observed in both field and laboratory collected populations as well as in insects reared on different diets ([Broderick et al., 2004](#); [Xiang et al., 2006](#); [Pinto-Tomás et al., 2011](#)), other studies reported no host specific resident communities that occurred, regardless of the insect diet ([Hammer et al., 2017](#)). It is possible that lepidopterans are less prone to forming robust “core” microbiomes due to several factors: (1) very high pH in the midgut, (2) low retention time of food, (3) lack of microbe housing structures in the intestinal tract, and (4) continual replacement of the peritrophic matrix ([Hammer et al., 2017](#)). Nevertheless, bacterial communities do continually associate with lepidopterans and influence a variety of important host processes ([Broderick, Raffa & Handelsman, 2006](#); [Anand et al., 2010](#); [Wang et al., 2017](#)).

Relatively few studies have assessed the *Spodoptera*-associated gut microbiome. In a recent study, the microbial diversity of *Spodoptera exigua* (Hübner) (Lepidoptera: Noctuidae) was examined by 16S rDNA sequence profiling ([Gao et al., 2019](#)). In *Spodoptera exigua*, the dominant bacterial clades are Proteobacteria and Firmicutes, with the predominant genus in larvae being *Enterococcus*. In *S. frugiperda*, previous studies have isolated several bacterial strains using culture-dependent methods ([De Almeida et al., 2017](#); [Acevedo et al., 2017](#)).

In this study, we used 16S rDNA sequence profiling to characterize the diversity of bacteria associated with populations of *S. frugiperda* in Kenya and assessed the prevalence of the *S. frugiperda* strains in these populations using mitochondrial COI gene sequences. Specifically, we characterized the structure of the circulating *S. frugiperda* populations in Kenya as well as the gut bacterial communities derived from both larval and adult specimens collected in different agro-ecological zones. Understanding pest population structures is important for understanding invasion patterns and planning with regards to strain-specific susceptibility of crops, whereas characterizing pest-associated microbiomes is a useful foundation for exploring insect-microbiome interactions that could be exploited to improve control strategies.

MATERIALS AND METHODS

Insect collection

Spodoptera frugiperda larvae were collected from infested maize fields in Kenya between June and December 2017 at the following locations: Ngeria (N00.37024 E035.9862) and Burnt Forest (N00.22505 E035.42479) in Uasin Gishu County; Msamia, Kitale (N00.98009 E034.97170) in Trans Nzoia County; Shimba Hills (S04.33228 E039.34361) in Kwale County and Chala Irrigation Scheme (S03.27338 E037.13816) and Wundanyi (S03.337538 E038.33612) in Taita Taveta County. Part of the field collected insects from each sampled region in Kenya were reared on fresh maize leaves in ventilated cages to pupation and eclosion at 27 °C and 60% humidity, while the rest were stored in absolute ethanol at –20 °C. We profiled the bacterial microbiome for 18 samples from four of these locations, whereas we included samples from all the sampled locations for mtDNA haplotyping (Fig. 1).

DNA extraction and 16S rDNA sequencing

Guts from nine live stage 5–6 larvae and nine 1 day old emerging adults from the Kenya collected samples were dissected separately in phosphate buffered saline (PBS) following surface sterilization and used for DNA extraction. Insects were surface sterilized in 70% ethanol, in 5% v/v sodium hypochlorite solution followed by three washes in PBS for 3 min in each solution. Each dissected gut tissue was homogenized in PBS using five 4 mm diameter ceramic beads in a two ml microfuge tube, using a TissueLyser II beadmill (Qiagen, Hilden, Germany). DNA was extracted using the ISOLATE II Genomic DNA Kit (Bioline, London, UK) according to the manufacturer's instructions. DNA extracted from gut samples was submitted for high throughput sequencing targeting the v4 region of the bacterial 16s rRNA gene using the Illumina Miseq platform (Center for Integrated Genomics, University of Lausanne, Switzerland). Sequence reads were checked for quality using FastQC v 0.11.28 (Andrews, 2010) and pre-processed to remove adapters and sequencing primers using Cutadapt v1.18 (Martin, 2011). Forward and reverse reads were imported into QIIME2-2018.11 (Boyle et al., 2018). The deblur plugin (Amir et al., 2017) was used to further filter the reads based on per base quality scores and merge the paired-end-reads and cluster reads into operational taxonomic units (OTUs). Taxonomic assignment was done using the blast classifier against the Silva132 reference database (Quast et al., 2013) at a 99% identity cutoff. OTU prevalence and variance based filtering as well as alpha and beta diversity measures were applied to the data in the Microbial Analyst Marker Data Profiling (Dhariwal et al., 2017). Shannon diversity indices were applied along with Mann–Whitney and analysis of variance statistics in profiling alpha diversity between sets of samples. Beta diversity was evaluated using Bray–Curtis and unweighted Unifrac distances. Significance testing was done using permutational multivariate analysis of variance (PERMANOVA) and visualization done through non-metric multidimensional scaling (NMDS) ordination. The empirical analysis of digital gene expression data in R (edgeR) algorithm (Robinson, McCarthy & Smyth, 2009) was used to evaluate differential abundance of bacterial genera reads between sample groups.




Figure 1 Sites from which *Spodoptera frugiperda* larvae were collected in Kenya.

Full-size  DOI: [10.7717/peerj.8701/fig-1](https://doi.org/10.7717/peerj.8701/fig-1)

All sequence reads were archived in the Sequence Read Archive under the BioProject: [PRJNA521837](https://www.ncbi.nlm.nih.gov/bioproject/PRJNA521837).

mtDNA haplotyping

DNA was extracted from surface-sterilized whole insects using the ISOLATE II Genomic DNA Kit (Bioline, London, UK) according to the manufacturer's instructions. Mitochondrial COI gene sequences were amplified from insect DNA by PCR using the primer LCO1490 and HCO2198 (Folmer *et al.*, 1994). Reactions were set up in total volumes of 10 μ l each, containing 5 \times MyTaq reaction buffer (5 mM dNTPs, 15 mM MgCl₂, stabilizers and enhancers) (Bioline, London, UK), 2 μ M of each primer, 0.25 mM MgCl₂ (Thermo Fischer Scientific, Waltham, MA, USA), 0.125 μ l MyTaq DNA




Figure 2 Composition of bacterial OTUs at Order level in screened *Spodoptera frugiperda* samples. Samples are denoted as “site name-developmental stage/sex_sample number”, where: l, larvae; m, male and f, female. [Full-size !\[\]\(1679558f37f6db0dd8360a2a7e913e90_img.jpg\) DOI: 10.7717/peerj.8701/fig-2](https://doi.org/10.7717/peerj.8701/fig-2)

polymerase (Bioline, London, UK), and 7.5 ng/μl of DNA template. These reactions were set up in a Master cycler Nexus gradient thermo-cycler (Thermo Fischer Scientific, Waltham, MA, USA) using the following cycling conditions: initial denaturation for 2 min at 95 °C, followed by 40 cycles of 30 s at 95 °C, 45 s at 50.6 °C and 1 min at 72 °C, then a final elongation step of 10 min at 72 °C. PCR products were separated by 1% agarose gel electrophoresis and visualized by ethidium bromide staining and UV trans-illumination. Direct sequencing was done for all host mtCOI gene and the sequences deposited in the GenBank.

RESULTS

A total of 457,501 sequence reads were retained after removal of spurious reads and all reads shorter than 220, which was the median length of all sequences with a quality score higher than 20. These sequences clustered into 1,796 OTUs. Of these, 197 OTUs survived low count and interquartile range-based variance filtering to eliminate OTUs that could arise from sequencing errors and contamination. OTUs initially characterized as “*Candidatus hamiltonella*” by comparison to the Silva132 reference database were re-analyzed by homology searches against the NCBI nr nucleotide database through blast (Altschul et al., 1990) and found to be *Pseudomonas*, highlighting a potential incorrect assignment in the reference database.

The most abundant bacterial Phyla observed across the FAW gut samples were Proteobacteria, Firmicutes, Bacteroidetes and a small proportion of Actinobacteria (Fig. S1). OTUs clustering in the orders Enterobacteriales and Pseudomonadales were predominant in the majority of the samples (Fig. 2).

We noticed that despite the high genus-level diversity between samples (Fig. 3), there were some similarities based on developmental stage and location. For example, there was a very high proportion of: (1) *Pseudomonas* in the two adult male samples from Chala, (2) *Citrobacter* in two larval samples from Kwale, (3) *Lysinibacillus* in two male samples




Figure 3 Genus level composition of (% of OTUs) in the different samples of *S. frugiperda*. Samples are denoted as “site name-developmental stage/sex_sample number”, where: l, larvae; m, male and f, female. The relative abundances of the 19 most abundant genera (with more than 10 counts after low count and variance filtering) are represented here. [Full-size !\[\]\(fd7fe780e8fd8eece60268c87d0c3e04_img.jpg\) DOI: 10.7717/peerj.8701/fig-3](https://doi.org/10.7717/peerj.8701/fig-3)

from Kitale and (4) *Enterococcus* in two larval samples from Ngeria. It was noted that *Stenotrophomonas*, *Sphingobacterium*, *Serratia*, *Pseudomonas*, *Morganella*, *Enterococcus* and *Delftia* were detected in both larvae and adult samples.

In one of the larval samples from the Ngeria site (Ngeria-l2), we observed an excessive number of non-bacterial reads. Through homology searches against the NCBI nr nucleotide database, these were found to be closely related to *Metarhizium rileyi* (Farl.) [Kepler et al. \(2014\)](#) (formerly *Nomuraea rileyi*), an entomopathogenic fungus that is, known to infect *S. frugiperda* ([Fig. 4](#)).

The bacterial OTU richness appeared to be higher in *S. frugiperda* larvae than adults, however this difference was not statistically significant (p -value: 0.062526; (Mann–Whitney) statistic: 19) using Shannon diversity metrics ([Fig. 5A](#)). In addition, no significant variation in OTU richness and abundance was observed between larvae from different sampling sites (p -value: 0.32834; (ANOVA) F -value: 1.3486) ([Fig. 5B](#)).

The composition of bacterial OTUs between larvae and adult *S. frugiperda* was observed to overlap although some significant dissimilarity ((PERMANOVA) F -value: 2.734; R -squared: 0.26715; p -value < 0.001 (NMDS) Stress = 0.13859) was recorded ([Fig. 6](#)). Similarly, OTU composition was observed to vary significantly among larval samples from




Figure 4 Neighbor-joining tree of fungus OTU detected in *S. frugiperda* sample (Nigeria-I2; in bold) and GenBank accessions of small subunit ribosomal RNA gene sequences from related fungi. Sequences are labeled by their GenBank accessions followed by genus, species and strain where available. Bootstrap values are indicated above the branches. Branches with a bootstrap value less than 50 are collapsed. A sequence from a species in the genus *Rozella* is included as an out-group.

Full-size DOI: [10.7717/peerj.8701/fig-4](https://doi.org/10.7717/peerj.8701/fig-4)

different sites ((PERMANOVA) F -value: 1.7511; R -squared: 0.36856; p -value < 0.037 (NMDS) Stress = 0.057109) (Fig. 7).

A significant differential abundance was observed for three bacterial genera between larvae and adult *S. frugiperda* samples using the EdgeR algorithm at an adjusted p -value of 0.05. Two of these: *Citrobacter* ($\log_2FC = 4.4178$, p value = $3.6E-6$, FDR = $7.218E-5$) and *Sphingobacterium* ($\log_2FC = 3.625$, p value = $1.01E-4$, FDR = 0.0010118) were more abundant in larvae whereas the third: *Lysinibacillus* ($\log_2FC = -3.2247$, p value = $4.4E-3$, FDR = 0.029375) was more abundant in adults (Fig. 8).

Based on mtDNA sequences, the *S. frugiperda* strains detected in this study were identical to strains from Canada, USA and Brazil, as well as strains that were recently




Figure 5 A comparison of the Shannon diversity indices for: (A) adult and larval samples from all sites and (B) larvae collected from different sites. The Shannon diversity index (H') was calculated based on the OTU-level of classification. The boxplots show the distribution of H' values across all samples. [Full-size](#) DOI: 10.7717/peerj.8701/fig-5




Figure 6 Non-metric multidimensional scaling (NMDS) ordination based on Bray-Curtis dissimilarities in bacterial communities detected in the *S. frugiperda* samples. Samples are colored according to their developmental stage and sex as indicated on the legend where: F, female; L, larvae and M, male. [Full-size](#) DOI: 10.7717/peerj.8701/fig-6




Figure 7 Non-metric multidimensional scaling (NMDS) ordination based on Weighted Unifrac distance dissimilarities in bacterial OTU composition between larval sample pairs from different sites. Samples are labeled by their collection site identifiers. [Full-size DOI: 10.7717/peerj.8701/fig-7](https://doi.org/10.7717/peerj.8701/fig-7)




Figure 8 Comparative abundance of (A) *Citrobacter*, (B) *Lysinibacillus* and (C) *Sphingobacterium* between adults and larvae of *S. frugiperda*. Abundance is shown on a log transformed scale of original counts. [Full-size DOI: 10.7717/peerj.8701/fig-8](https://doi.org/10.7717/peerj.8701/fig-8)

reported in Kenya and parts of Africa and India (Fig. 9). All the samples clustered in two major clades widely referred to as either the “Rice” or the “Corn” strain (hereafter referred to as R-strain and C-strain). We investigated the frequency of mtDNA haplotypes of




Figure 9 Neighbor-joining tree based on mtCOI sequences of *S. frugiperda* from the GenBank and representative haplotypes from this study (in bold). Bootstrap values are indicated above branches. Branches with bootstrap values less than 50 are collapsed. A sequence from *Spodoptera exigua* is included as an out-group. Sequences are labeled with their GenBank accession numbers, collection site where available and country of collection. [Full-size !\[\]\(e4c51d9db35ee9651ed60d72acdb782c_img.jpg\) DOI: 10.7717/peerj.8701/fig-9](https://doi.org/10.7717/peerj.8701/fig-9)




Figure 10 Mitochondrial COI haplotype map of the *S. frugiperda* samples collected at various sites in Kenya. Node size is proportional to number of samples and individual samples are represented as fractions of the nodes. A sequence of *Spoladea recurvalis* is included as an out-group. Sequences for all samples are accessible from GenBank using the accessions: MK492929–MK493010.

Full-size  DOI: 10.7717/peerj.8701/fig-10

S. frugiperda samples collected at several sites in Kenya. Overall, 90% of the samples ($n = 85$) clustered as R-strain, whereas 10% ($n = 9$) clustered as C-strain. Proportions of the R-strain in populations at the different sites were 100% ($n = 6$) for Burnt Forest, 83% ($n = 6$) for Chala, 86% ($n = 7$) for Wundanyi, 82% ($n = 11$) for Kitale, 91% ($n = 35$) for Kwale and 82% ($n = 17$) for Ngeria (Fig. 10).

DISCUSSION

We found that the gut bacterial communities of most *S. frugiperda* samples were dominated by Proteobacteria. This observation is similar to proportions reported in other phytophagous insects, in particular lepidopterans (Belda et al., 2011; Xia et al., 2013, 2017; Landry et al., 2015; Ramya et al., 2016; Snyman et al., 2016; Strano et al., 2018; Chen et al., 2018). Only three samples, two adult males from Kitale (Kitale-m2 and Kitale-m3) and one larva from Ngeria (Ngeria-l2) were dominated by Firmicutes. The four genera of bacteria, *Pseudomonas*, *Delftia*, *Enterococcus* and *Serratia*, that were recorded in this study have previously been isolated from *S. frugiperda* (De Almeida et al., 2017; Acevedo et al., 2017). Surprisingly, *Staphylococcus*, *Microbacterium*, *Arthrobacter* and *Leclercia* that were previously isolated from *S. frugiperda* in Brazil (De Almeida et al., 2017) were not found in any of the samples we profiled in Kenya.

Similarly, *Pantoea*, *Enterobacter*, *Raoultella* and *Klebsiella* previously identified in oral secretions of *S. frugiperda* in Pennsylvania, USA (Acevedo et al., 2017) were not found in the profiled Kenyan samples. Six of the detected bacterial genera: *Enterococcus*, *Pseudomonas*, *Chryseobacterium*, *Sphingobacterium*, *Ochrobactrum* and *Acinetobacter*, have been detected using a similar sequencing approach in both *S. frugiperda* as well as in the corn earworm *Helicoverpa zea* (Jones et al., 2019). Similarly, seven of the detected

bacterial genera: *Enterococcus*, *Pseudomonas*, *Comamonas*, *Stenotrophomonas*, *Eshcherichia-Shigella*, *Acinetobacter* and *Carnobacterium*, have been reported using a similar approach in the beet armyworm, *S. exigua* (Gao *et al.*, 2019). This suggests that some bacterial genera often associate with lepidopteran insects, although it is difficult to define a core microbiota for such a diverse insect order. The OTUs classified as *Candidiatus hamiltonella* using the Silva database were further investigated and reclassified as *Pseudomonas*. *Candidiatus hamiltonella* has been recorded in whiteflies, psyllids and phloem-feeding relatives of the aphids (Russell & Moran, 2005) but not among lepidopteran insects.

We observed significant differences in OTU composition between larvae from different sites. This was most likely caused by complex biological and environmental factors in the diverse agro-ecological zones that were sampled. Diet is known to strongly influence the microbiome of lepidopterans (Strano *et al.*, 2018; Sittenfeld *et al.*, 2002; Priya *et al.*, 2012; Montagna *et al.*, 2016), however in this study all samples were collected from maize plants. Hence, the observed compositional differences are not likely to be caused solely by diet. It is interesting that many of the detected bacterial genera such as *Stenotrophomonas*, *Sphingobacterium*, *Serratia*, *Pseudomonas*, *Morganella*, *Enterococcus* and *Delftia* were found in both life stages, which suggests that gut bacterial community members are transmitted across developmental stages. Bacteria that are continually transmitted across developmental stages (and across generations) may evolve a closer, mutualistic relationship with their hosts (Moran, 2006). Future studies should investigate the effects of these microbes on host fitness and investigate the extent to which they are vertically transmitted from parents to offspring. In contrast, *Citrobacter* and *Sphingobacterium* were observed to be differentially abundant in larvae than in adults, a likely indicator that these two genera may be part of the fraction of bacterial communities that are lost during transition of *S. frugiperda* into the adult stage. *Lysinibacillus*, on the other hand, was more abundant in adults than in larvae and therefore could have an adult-specific function.

Notably, we identified *Serratia*, *Lysinibacillus* (formerly *Bacillus*) and *Pseudomonas*, species of which have been reported to have entomopathogenic properties (Castagnola & Stock, 2014). In addition, one sample had a high number of reads attributed to a relative of a fungal entomopathogen, *Metarhizium rileyi*, which previously has been isolated and tested for efficacy against *S. frugiperda* (Maniania & Fargues, 1985; Mallapur *et al.*, 2018). However, there was no record of the use of any fungal biopesticides in any of the sampled sites. This detection could however be an indication that *S. frugiperda* might be refractory to the fungus, in which case such a fungus could be exploited in alternative control techniques such as in the delivery of dsRNA. It may be worthwhile to re-examine the pathogenicity of these microbes for *S. frugiperda* and to determine if they could be incorporated into biological pest management strategies (Ruiu, 2015).

Based on the mtCOI gene sequences, we observed two mtDNA haplotypes in Kenya (C- and R-strains), despite the fact that all of these insects were obtained from maize. These findings confirm that both haplotypes are present in Kenya, as has been demonstrated for other countries in Africa (Rwomushana *et al.*, 2018). The majority of the

S. frugiperda samples collected were characterized as R-strain suggesting that this strain is dominant in *S. frugiperda* populations in Kenya. These observations are in agreement with a previous study (Nagoshi *et al.*, 2018) that observed that C- and R-strains appear to have an East-West axis alignment in the African region, with Eastern Africa having progressively lower frequencies of the C-strain (Nagoshi *et al.*, 2018). We noted that some variants of the R-strain have been reported in other places such as Ghana and India, but those variants were not detected in this study. It is interesting to note that in addition to an R-strain similar to the one detected in Kenya, a variant differing by a single nucleotide polymorphism in the sequenced region of the mtCOI gene has been recorded from various locations in India. This variant has however not been reported in Africa. It is therefore possible that the invasion into India may not have come directly from the African continent, or invasion could have included strains from Africa and elsewhere.

CONCLUSIONS

We characterized the gut bacterial communities in *S. frugiperda* larvae and adult samples collected from several locations in Kenya, finding some important differences and similarities across samples and in relation to other studies on this species (Acevedo *et al.*, 2017; De Almeida *et al.*, 2017). Characterizing the gut microbial symbionts of this pest species in Africa can be seen as an important first step towards the development of novel, cost-effective symbiont and entomopathogen-based control strategies. In addition, the population structure of this pest in Kenya was investigated. Understanding the population structure, dynamics and bio-ecology of invasive species is important for identifying their invasion patterns and for informing cropping systems especially where pest species compositions associate with different host plant usage.

ACKNOWLEDGEMENTS

The authors acknowledge the technical support of Mr. Peter Malusi during sampling.

ADDITIONAL INFORMATION AND DECLARATIONS

Funding

Financial support for this research came from the following organizations and agencies: European Union Funded project “Integrated pest management” strategy to counter the threat of invasive fall armyworm to food security in Eastern Africa (FAW-IPM) (FOOD/2018/402-634); UK Aid from the UK Government; Swedish International Development Cooperation Agency (Sida); the Swiss Agency for Development and Cooperation (SDC); Federal Democratic Republic of Ethiopia; and the Kenyan Government. Financial assistance for sequencing at the Center for Integrated Genomics, Lausanne, was provided by the Global Health Institute of the École Polytechnique Fédérale de Lausanne. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Grant Disclosures

The following grant information was disclosed by the authors:

European Union: FOOD/2018/402-634.

UK Aid from the UK Government.

Swedish International Development Cooperation Agency (Sida).

Swiss Agency for Development and Cooperation (SDC).

Federal Democratic Republic of Ethiopia.

Kenyan Government.

Global Health Institute of the École Polytechnique Fédérale de Lausanne.

Competing Interests

The authors declare that they have no competing interests.

Author Contributions

- Joseph Gichuhi performed the experiments, analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the paper, and approved the final draft.
- Subramanian Sevgan conceived and designed the experiments, authored or reviewed drafts of the paper, and approved the final draft.
- Fathiya Khamis conceived and designed the experiments, authored or reviewed drafts of the paper, and approved the final draft.
- Johnnie Van den Berg conceived and designed the experiments, authored or reviewed drafts of the paper, and approved the final draft.
- Hannalene du Plessis conceived and designed the experiments, authored or reviewed drafts of the paper, and approved the final draft.
- Sunday Ekesi conceived and designed the experiments, authored or reviewed drafts of the paper, and approved the final draft.
- Jeremy K. Herren conceived and designed the experiments, analyzed the data, authored or reviewed drafts of the paper, and approved the final draft.

Data Availability

The following information was supplied regarding data availability:

Gut bacteria reads are available in the Sequence Read Archive (SRA) BioProject:

[PRJNA521837](https://www.ncbi.nlm.nih.gov/bioproject/PRJNA521837).

Fall armyworm mtCOI sequences are available at GenBank: [MK492929](https://www.ncbi.nlm.nih.gov/genbank/MK492929) to [MK493010](https://www.ncbi.nlm.nih.gov/genbank/MK493010).

Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.8701#supplemental-information>.

REFERENCES

- Acevedo FE, Peiffer M, Tan C-W, Stanley BA, Stanley A, Wang J, Jones AG, Hoover K, Rosa C, Luthe D, Felton G. 2017. Fall armyworm-associated gut bacteria modulate plant defense responses. *Molecular Plant–Microbe Interactions* **30**(2):127–137
[DOI 10.1094/MPMI-11-16-0240-R](https://doi.org/10.1094/MPMI-11-16-0240-R).

- Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic local alignment search tool. *Journal of Molecular Biology* 215(3):403–410 DOI 10.1016/S0022-2836(05)80360-2.
- Amir A, McDonald D, Navas-Molina JA, Kopylova E, Morton JT, Xu ZZ, Kightley EP, Thompson LR, Hyde ER, Gonzalez A, Knight R. 2017. Deblur rapidly resolves single-nucleotide community sequence patterns. *mSystems* 2:1–7.
- Anand AAP, Vennison SJ, Sankar SG, Prabhu DIG, Vasan PT, Raghuraman T, Geoffrey CJ, Vendan SE. 2010. Isolation and characterization of bacteria from the gut of *Bombyx mori* that degrade cellulose, xylan, pectin and starch and their impact on digestion. *Journal of Insect Science* 10(107):1–20 DOI 10.1673/031.010.10701.
- Andrews S. 2010. FastQC: a quality control tool for high throughput sequence data. Available at <https://www.bibsonomy.org/bibtex/2b6052877491828ab53d3449be9b293b3/ozborn>.
- Arora AK, Douglas AE. 2017. Hype or opportunity? Using microbial symbionts in novel strategies for insect pest control. *Journal of Insect Physiology* 103:10–17 DOI 10.1016/j.jinsphys.2017.09.011.
- Banerjee R, Hasler J, Meagher R, Nagoshi R, Hietala L, Huang F, Narva K, Jurat-Fuentes JL. 2017. Mechanism and DNA-based detection of field-evolved resistance to transgenic Bt corn in fall armyworm (*Spodoptera frugiperda*). *Scientific Reports* 7(1):1–10 DOI 10.1038/s41598-016-0028-x.
- Beck JJ, Vannette RL. 2017. Harnessing insect-microbe chemical communications to control insect pests of agricultural systems. *Journal of Agricultural and Food Chemistry* 65(1):23–28 DOI 10.1021/acs.jafc.6b04298.
- Belda E, Pedrola L, Peretó J, Martínez-Blanch JF, Montagud A, Navarro E, Urchueguía J, Ramón D, Moya A, Porcar M. 2011. Microbial diversity in the midguts of field and lab-reared populations of the European corn borer *Ostrinia nubilalis*. *PLOS ONE* 6(6):e21751 DOI 10.1371/journal.pone.0021751.
- Biere A, Bennett AE. 2013. Three-way interactions between plants, microbes and insects. *Functional Ecology* 27(3):567–573 DOI 10.1111/1365-2435.12100.
- Botha AS, Erasmus A, du Plessis H, Van den Berg J. 2019. Efficacy of Bt maize for control of *Spodoptera frugiperda* (Lepidoptera: Noctuidae) in South Africa. *Journal of Economic Entomology* 112(3):1260–1266 DOI 10.1093/jee/toz048.
- Boylen E, Rideout JR, Dillon MR, Bokulich NA, Abnet C, Al-Ghalith GA, Alexander H, Alm EJ, Arumugam M, Asnicar F, Bai Y, Bisanz JE, Bittinger K, Brejnrod A, Brislawn CJ, Brown CT, Callahan BJ, Caraballo-Rodríguez AM, Chase J, Cope E, Da Silva R, Dorrestein PC, Douglas GM, Durall DM, Duvallet C, Edwardson CF, Ernst M, Estaki M, Fouquier J, Gauglitz JM, Gibson DL, Gonzalez A, Gorlick K, Guo J, Hillmann B, Holmes S, Holste H, Huttenhower C, Huttley GA, Kang KB, Janssen S, Jarmusch AK, Jiang L, Kaehler B, Keefe CR, Keim P, Kelley ST, Knights D, Koester I, Kosciolk T, Kreps J, Langille MGI, Lee J, Ley R, Liu Y-X, Lofffield E, Lozupone C, Maher M, Marotz C, Martin BD, McDonald D, McIver LJ, Melnik AV, Metcalf JL, Morgan SC, Morton J, Naimey AT, Navas-Molina JA, Nothias LF, Orchanian SB, Pearson T, Peoples SL, Petras D, Preuss ML, Pruesse E, Rasmussen LB, Rivers A, Robeson MS II, Rosenthal P, Segata N, Shaffer M, Shiffer A, Sinha R, Song SJ, Spear JR, Swafford AD, Thompson LR, Torres PJ, Trinh P, Tripathi A, Turnbaugh PJ, Ul-Hasan S, Van Der Hooft JJJ, Vargas F, Vázquez-Baeza Y, Vogtmann E, Von Hippel M, Walters W, Wan Y, Wang M, Warren J, Weber KC, Williamson CHD, Willis AD, Xu ZZ, Zaneveld JR, Zhang Y, Zhu Q, Knight R, Caporaso JG. 2018. QIIME 2: reproducible, interactive, scalable, and extensible microbiome data science. *PeerJ Preprints*. 6:e27295v2 DOI 10.7287/peerj.preprints.27295.

- Brady CM, White JA. 2013.** Cowpea aphid (*Aphis craccivora*) associated with different host plants has different facultative endosymbionts. *Ecological Entomology* **38**(4):433–437 DOI [10.1111/een.12020](https://doi.org/10.1111/een.12020).
- Broderick NA, Raffa KF, Goodman RM, Handelsman J. 2004.** Census of the bacterial community of the gypsy moth larval midgut by using culturing and culture-independent methods. *Applied and Environmental Microbiology* **70**(1):293–300 DOI [10.1128/AEM.70.1.293-300.2004](https://doi.org/10.1128/AEM.70.1.293-300.2004).
- Broderick NA, Raffa KF, Handelsman J. 2006.** Midgut bacteria required for *Bacillus thuringiensis* insecticidal activity. *Proceedings of the National Academy of Sciences of the United States of America* **103**(41):15196–15199 DOI [10.1073/pnas.0604865103](https://doi.org/10.1073/pnas.0604865103).
- Castagnola A, Stock SP. 2014.** Common virulence factors and tissue targets of entomopathogenic bacteria for biological control of Lepidopteran pests. *Insects* **5**(1):139–166 DOI [10.3390/insects5010139](https://doi.org/10.3390/insects5010139).
- Chen B, Du K, Sun C, Vimalanathan A, Liang X, Li Y, Wang B, Lu X, Li L, Shao Y. 2018.** Gut bacterial and fungal communities of the domesticated silkworm (*Bombyx mori*) and wild mulberry-feeding relatives. *ISME Journal* **12**(9):2252–2262 DOI [10.1038/s41396-018-0174-1](https://doi.org/10.1038/s41396-018-0174-1).
- Crotti E, Balloi A, Hamdi C, Sansonno L, Marzorati M, Gonella E, Favia G, Cherif A, Bandi C, Alma A, Daffonchio D. 2012.** Microbial symbionts: a resource for the management of insect-related problems. *Microbial Biotechnology* **5**(3):307–317 DOI [10.1111/j.1751-7915.2011.00312.x](https://doi.org/10.1111/j.1751-7915.2011.00312.x).
- Day R, Abrahams P, Bateman M, Beale T, Clotey V, Cock M, Colmenarez Y, Corniani N, Early R, Godwin J, Gomez J, Moreno PG, Murphy ST, Oppong-Mensah B, Phiri N, Pratt C, Silvestri S, Witt A. 2017.** Fall armyworm: impacts and implications for Africa. *Outlooks on Pest Management* **28**(5):196–201 DOI [10.1564/v28_oct_02](https://doi.org/10.1564/v28_oct_02).
- De Almeida LG, De Moraes LAB, Trigo JR, Omoto C, Cônsoli FL. 2017.** The gut microbiota of insecticide-resistant insects houses insecticide-degrading bacteria: a potential source for biotechnological exploitation. *PLOS ONE* **12**(3):e0174754 DOI [10.1371/journal.pone.0174754](https://doi.org/10.1371/journal.pone.0174754).
- Deole S, Paul N. 2018.** First report of fall army worm, *Spodoptera frugiperda* (J.E. Smith), their nature of damage and biology on maize crop at Raipur. *Chhattisgarh Journal of Entomology and Zoology Studies* **6**:219–221.
- Dhariwal A, Chong J, Habib S, King IL, Agellon LB, Xia J. 2017.** MicrobiomeAnalyst: a web-based tool for comprehensive statistical, visual and meta-analysis of microbiome data. *Nucleic Acids Research* **45**(W1):W180–W188 DOI [10.1093/nar/gkx295](https://doi.org/10.1093/nar/gkx295).
- Douglas AE. 2009.** The microbial dimension in insect nutritional ecology. *Functional Ecology* **23**(1):38–47 DOI [10.1111/j.1365-2435.2008.01442.x](https://doi.org/10.1111/j.1365-2435.2008.01442.x).
- Douglas AE. 2015.** Multiorganismal insects: diversity and function of resident microorganisms. *Annual Review of Entomology* **60**(1):17–34 DOI [10.1146/annurev-ento-010814-020822](https://doi.org/10.1146/annurev-ento-010814-020822).
- FAO. 2018.** *Sustainable management of the fall armyworm in Africa: a framework for partnership*. Rome: Food and Agriculture Organization of the United Nations.
- Ferrari J, Vavre F, Lyon D. 2011.** Bacterial symbionts in insects or the story of communities affecting communities. *Philosophical Transactions of the Royal Society B: Biological Sciences* **366**(1569):1389–1400 DOI [10.1098/rstb.2010.0226](https://doi.org/10.1098/rstb.2010.0226).
- Flagel L, Lee YW, Wanjugi H, Swarup S, Brown A, Wang J, Kraft E, Greenplate J, Simmons J, Adams N, Wang Y, Martinelli S, Haas JA, Gowda A, Head G. 2018.** Mutational disruption of the ABCC2 gene in fall armyworm, *Spodoptera frugiperda*, confers resistance to the Cry1Fa and Cry1A.105 insecticidal proteins. *Scientific Reports* **8**:7255 DOI [10.1038/s41598-018-25491-9](https://doi.org/10.1038/s41598-018-25491-9).

- Folmer O, Black M, Hoeh W, Lutz R, Vrijenhoek R. 1994.** DNA primers for amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. *Molecular Marine Biology and Biotechnology* **3(10)**:294–299.
- Frago E, Dicke M, Godfray HCJ. 2012.** Insect symbionts as hidden players in insect–plant interactions. *Trends in Ecology & Evolution* **27(12)**:705–711 DOI [10.1016/j.tree.2012.08.013](https://doi.org/10.1016/j.tree.2012.08.013).
- Gao X, Li W, Luo J, Zhang L, Ji J, Zhu X, Wang L, Zhang S, Cui J. 2019.** Biodiversity of the microbiota in *Spodoptera exigua* (Lepidoptera: Noctuidae). *Journal of Applied Microbiology* **126(4)**:1199–1208 DOI [10.1111/jam.14190](https://doi.org/10.1111/jam.14190).
- Garcia ES, Castro DP, Figueiredo MB, Azambuja P. 2010.** Immune homeostasis to microorganisms in the guts of triatomines (Reduviidae): a review. *Memorias do Instituto Oswaldo Cruz* **105(5)**:605–610 DOI [10.1590/S0074-02762010000500001](https://doi.org/10.1590/S0074-02762010000500001).
- Goergen G, Kumar PL, Sankung SB, Togola A, Tamò M. 2016.** First report of outbreaks of the fall armyworm *Spodoptera frugiperda* (J.E. Smith) (Lepidoptera, Noctuidae), a new alien invasive pest in West and Central Africa. *PLOS ONE* **11**:1–9 DOI [10.1371/journal.pone.0165632](https://doi.org/10.1371/journal.pone.0165632).
- Hailu G, Niassy S, Zeyaur KR, Ochatum N, Subramanian S. 2018.** Maize-legume intercropping and push-pull for management of fall armyworm, stemborers, and striga in Uganda. *Agronomy Journal* **110(6)**:2513–2522 DOI [10.2134/agronj2018.02.0110](https://doi.org/10.2134/agronj2018.02.0110).
- Hammer TJ, Janzen H, Hallwachs D, Jaffe W, Fierera SP, Fierer N. 2017.** Caterpillars lack a resident gut microbiome. *Proceedings of the National Academy of Sciences of the United States of America* **114(36)**:9641–9646 DOI [10.1073/pnas.1707186114](https://doi.org/10.1073/pnas.1707186114).
- Huber DM, Hugh-Jones ME, Rust MK, Sheffield SR, Simberloff D, Taylor CR. 2002.** Invasive pest species: impacts on agricultural production, natural resources and the environment. *Council for Agricultural Science and Technology* **20**:1–18.
- Jakka SRK, Gong L, Hasler J, Banerjee R, Sheets JJ, Narva K, Blanco CA, Jurat-Fuentes JL. 2016.** Field-evolved mode 1 resistance of the fall armyworm to transgenic Cry1Fa-expressing corn associated with reduced Cry1Fa toxin binding and midgut alkaline phosphatase expression. *Applied and Environmental Microbiology* **82(4)**:1023–1034 DOI [10.1128/AEM.02871-15](https://doi.org/10.1128/AEM.02871-15).
- Jones AG, Mason CJ, Felton GW, Hoover K. 2019.** Host plant and population source drive diversity of microbial gut communities in two polyphagous insects. *Scientific Reports* **9(1)**:2792 DOI [10.1038/s41598-019-39163-9](https://doi.org/10.1038/s41598-019-39163-9).
- Kepler RM, Humber RA, Bischoff JF, Rehner SA. 2014.** Clarification of generic and species boundaries for *Metarhizium* and related fungi through multigene phylogenetics. *Mycologia* **106(4)**:811–829.
- Kikuchi Y, Hayatsu M, Hosokawa T, Nagayama A, Tago K, Fukatsu T. 2012.** Symbiont-mediated insecticide resistance. *Proceedings of the National Academy of Sciences of the United States of America* **109(22)**:8618–8622 DOI [10.1073/pnas.1200231109](https://doi.org/10.1073/pnas.1200231109).
- Landry M, Comeau AM, Derome N, Cusson M, Levesque RC. 2015.** Composition of the spruce budworm (*Choristoneura fumiferana*) midgut microbiota as affected by rearing conditions. *PLOS ONE* **10(12)**:e0144077 DOI [10.1371/journal.pone.0144077](https://doi.org/10.1371/journal.pone.0144077).
- Mallapur CP, Naik AK, Hagari S, Praveen T, Patil RK. 2018.** Potentiality of *Nomuraea rileyi* (Farlow) Samson against the fall armyworm, *Spodoptera frugiperda* (J.E. Smith) infesting maize. *Journal of Entomology and Zoology Studies* **6**:1062–1067.
- Maniania NK, Fargues J. 1985.** Susceptibility of the fall armyworm, *Spodoptera frugiperda*, to the fungal pathogens *Paecilomyces fumosoroseus* and *Nomuraea rileyi*. *Florida Entomologist* **68(1)**:178–183 DOI [10.2307/3494343](https://doi.org/10.2307/3494343).
- Martin M. 2011.** Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet journal* **17(1)**:10 DOI [10.14806/ej.17.1.200](https://doi.org/10.14806/ej.17.1.200).

- McLean AHC, Godfray HCJ. 2015.** Evidence for specificity in symbiont-conferred protection against parasitoids. *Proceedings of the Royal Society B: Biological Sciences* **282(1811)**:20150977 DOI [10.1098/rspb.2015.0977](https://doi.org/10.1098/rspb.2015.0977).
- Midega CAO, Pittchar JO, Pickett JA, Hailu GW, Khan ZR. 2018.** A climate-adapted push-pull system effectively controls fall armyworm, *Spodoptera frugiperda* (J.E. Smith), in maize in East Africa. *Crop Protection* **105**:10–15 DOI [10.1016/j.cropro.2017.11.003](https://doi.org/10.1016/j.cropro.2017.11.003).
- Montagna M, Mereghetti V, Gargari G, Guglielmetti S, Faoro F, Lozzia G, Locatelli D, Limonta L. 2016.** Evidence of a bacterial core in the stored products pest *Plodia interpunctella*: the influence of different diets. *Environmental Microbiology* **18(12)**:4961–4973 DOI [10.1111/1462-2920.13450](https://doi.org/10.1111/1462-2920.13450).
- Moran NA. 2006.** Symbiosis. *Current Biology* **16(20)**:R866–R871 DOI [10.1016/j.cub.2006.09.019](https://doi.org/10.1016/j.cub.2006.09.019).
- Nagoshi RN, Goergen G, Du Plessis H, Van Den Berg J, Meagher R Jr. 2019.** Genetic comparisons of fall armyworm populations from 11 countries spanning sub-Saharan Africa provide insights into strain composition and migratory behaviors. *Scientific Reports* **9(1)**:1–11 DOI [10.1038/s41598-019-44744-9](https://doi.org/10.1038/s41598-019-44744-9).
- Nagoshi RN, Goergen G, Tounou KA, Agboka K, Koffi D, Meagher RL. 2018.** Analysis of strain distribution, migratory potential, and invasion history of fall armyworm populations in northern Sub-Saharan Africa. *Scientific Reports* **8(1)**:3710 DOI [10.1038/s41598-018-21954-1](https://doi.org/10.1038/s41598-018-21954-1).
- Nagoshi RN, Meagher RL. 2004.** Behavior and distribution of the two fall armyworm host strains in Florida. *Florida Entomologist* **87(4)**:440–449 DOI [10.1653/0015-4040\(2004\)087\[0440:BADOTT\]2.0.CO;2](https://doi.org/10.1653/0015-4040(2004)087[0440:BADOTT]2.0.CO;2).
- Perilla-Henao LM, Casteel CL. 2016.** Vector-borne bacterial plant pathogens: interactions with Hemipteran insects and plants. *Frontiers in Plant Science* **7(1088)**:1163 DOI [10.3389/fpls.2016.01163](https://doi.org/10.3389/fpls.2016.01163).
- Pinto-Tomás AA, Sittenfeld A, Uribe-Lorío L, Chavarría F, Mora M, Janzen DH, Goodman RM, Simon HM. 2011.** Comparison of midgut bacterial diversity in tropical caterpillars (Lepidoptera: Saturniidae) fed on different diets. *Environmental Entomology* **40(5)**:1111–1122 DOI [10.1603/EN11083](https://doi.org/10.1603/EN11083).
- Priya NG, Ojha A, Kajla MK, Raj A, Rajagopal R. 2012.** Host plant induced variation in gut bacteria of *Helicoverpa armigera*. *PLOS ONE* **7(1)**:e30768 DOI [10.1371/journal.pone.0030768](https://doi.org/10.1371/journal.pone.0030768).
- Quast C, Prusse EP, Yilmaz P, Gerken J, Schweer T, Yarza P, Peplies J, Glockner FO. 2013.** The SILVA ribosomal RNA gene database project: improved data processing and web-based tools. *Nucleic Acids Research* **41(D1)**:D590–D596 DOI [10.1093/nar/gks1219](https://doi.org/10.1093/nar/gks1219).
- Ramya SL, Venkatesan T, Srinivasa Murthy KS, Jalali SK, Verghese A. 2016.** Detection of carboxylesterase and esterase activity in culturable gut bacterial flora isolated from diamondback moth, *Plutella xylostella* (Linnaeus), from India and its possible role in indoxacarb degradation. *Brazilian Journal of Microbiology* **47(2)**:327–336 DOI [10.1016/j.bjm.2016.01.012](https://doi.org/10.1016/j.bjm.2016.01.012).
- Robinson MD, McCarthy DJ, Smyth GK. 2009.** edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics* **26(1)**:139–140 DOI [10.1093/bioinformatics/btp616](https://doi.org/10.1093/bioinformatics/btp616).
- Ruiu L. 2015.** Insect pathogenic bacteria in integrated pest management. *Insects* **6(2)**:352–367 DOI [10.3390/insects6020352](https://doi.org/10.3390/insects6020352).
- Russell JA, Moran NA. 2005.** Horizontal transfer of bacterial symbionts: heritability and fitness effects in a novel aphid host. *Applied and Environmental Microbiology* **71(12)**:7987–7994 DOI [10.1128/AEM.71.12.7987-7994.2005](https://doi.org/10.1128/AEM.71.12.7987-7994.2005).
- Rwomushana I, Bateman M, Beale T, Beseh P, Cameron K, Chiluba M, Clotney V, Davis T, Day R, Early R, Godwin J, Gonzalez-Moreno P, Kansime M, Kenis M, Makale F, Idah M,**

- Murphy S, Nunda W, Phiri N, Pratt C, Tambo J. 2018.** *Fall armyworm: impacts and implications for Africa: evidence note update (October 2018)*. UK: CABI.
- Sisay B, Simiyu J, Malusi P, Likhayo P, Mendesil E, Elibariki N, Wakgari M, Ayalew G, Tefera T. 2018.** First report of the fall armyworm, *Spodoptera frugiperda* (Lepidoptera: Noctuidae), natural enemies from Africa. *Journal of Applied Entomology* **142(8)**:800–804 DOI [10.1111/jen.12534](https://doi.org/10.1111/jen.12534).
- Sisodiya DB, Raghunandan BL, Bhatt NA, Verma HS, Shewale CP, Timbadiya BG, Borad PK. 2018.** The fall armyworm, *Spodoptera frugiperda* (J.E. Smith) (Lepidoptera: Noctuidae); first report of new invasive pest in maize fields of Gujarat. *India Journal of Entomology and Zoology Studies* **6**:2089–2091.
- Sittenfeld A, Uribe-lorío L, Mora M, Nielsen V, Arrieta G, Janzen DH. 2002.** Does a polyphagous caterpillar have the same gut microbiota when feeding on different species of food plants? *Revista de Biología Tropical* **50**:547–560.
- Snyman M, Gupta AK, Bezuidenhout CC, Claassens S, Van den Berg J. 2016.** Gut microbiota of *Busseola fusca* (Lepidoptera: Noctuidae). *World Journal of Microbiology and Biotechnology* **32(7)**:115 DOI [10.1007/s11274-016-2066-8](https://doi.org/10.1007/s11274-016-2066-8).
- Strano CP, Malacrino A, Campolo O, Palmeri V. 2018.** Influence of host plant on *Thaumetopoea pityocampa* gut bacterial community. *Microbial Ecology* **75(2)**:487–494 DOI [10.1007/s00248-017-1019-6](https://doi.org/10.1007/s00248-017-1019-6).
- Su Q, Zhou X, Zhang Y. 2013.** Symbiont-mediated functions in insect hosts. *Communicative & Integrative Biology* **6(3)**:e23804 DOI [10.4161/cib.23804](https://doi.org/10.4161/cib.23804).
- Ubeda C, Djukovic A, Isaac S. 2017.** Roles of the intestinal microbiota in pathogen protection. *Clinical & Translational Immunology* **6(2)**:e128 DOI [10.1038/cti.2017.2](https://doi.org/10.1038/cti.2017.2).
- Wang J, Peiffer M, Hoover K, Rosa C, Zeng R, Felton GW. 2017.** *Helicoverpa zea* gut-associated bacteria indirectly induce defenses in tomato by triggering a salivary elicitor(s). *New Phytologist* **214(3)**:1294–1306 DOI [10.1111/nph.14429](https://doi.org/10.1111/nph.14429).
- Wei G, Lai Y, Wang G, Chen H, Li F, Wang S. 2017.** Insect pathogenic fungus interacts with the gut microbiota to accelerate mosquito mortality. *Proceedings of the National Academy of Sciences of the United States of America* **114(23)**:5994–5999 DOI [10.1073/pnas.1703546114](https://doi.org/10.1073/pnas.1703546114).
- Xia X, Gurr GM, Vasseur L, Zheng D, Zhong H, Qin B, Lin J, Wang Y, Song F, Li Y, Lin H, You M. 2017.** Metagenomic sequencing of diamondback moth gut microbiome unveils key holobiont adaptations for herbivory. *Frontiers in Microbiology* **8**:663 DOI [10.3389/fmicb.2017.00663](https://doi.org/10.3389/fmicb.2017.00663).
- Xia X, Sun B, Gurr GM, Vasseur L, Xue M, You M. 2018.** Gut microbiota mediate insecticide resistance in the diamondback moth, *Plutella xylostella* (L.). *Frontiers in Microbiology* **9**:1–10 DOI [10.3389/fmicb.2018.00025](https://doi.org/10.3389/fmicb.2018.00025).
- Xia X, Zheng D, Zhong H, Qin B, Gurr GM, Vasseur L, Lin H, Bai J, He W, You M. 2013.** DNA sequencing reveals the midgut microbiota of diamondback moth, *Plutella xylostella* (L.) and a possible relationship with insecticide resistance. *PLOS ONE* **8**:e68852 DOI [10.1371/journal.pone.0068852](https://doi.org/10.1371/journal.pone.0068852).
- Xiang H, Wei G-F, Jia S, Huang J, Miao X-X, Zhou Z, Zhao L-P, Huang Y-P. 2006.** Microbial communities in the larval midgut of laboratory and field populations of cotton bollworm (*Helicoverpa armigera*). *Canadian Journal of Microbiology* **52(11)**:1085–1092 DOI [10.1139/w06-064](https://doi.org/10.1139/w06-064).
- Yu SJ. 1991.** Insecticide resistance in the fall armyworm, *Spodoptera frugiperda* (J.E. Smith). *Pesticide Biochemistry and Physiology* **39(1)**:84–91 DOI [10.1016/0048-3575\(91\)90216-9](https://doi.org/10.1016/0048-3575(91)90216-9).